

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number  
**WO 02/04522 A2**

- (51) International Patent Classification<sup>7</sup>: **C07K 16/00**
- (21) International Application Number: **PCT/EP01/07878**
- (22) International Filing Date: **9 July 2001 (09.07.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:  
**09/613,017 10 July 2000 (10.07.2000) US**
- (71) Applicant (for all designated States except AT, US): **NOVARTIS AG [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).**
- (71) Applicant (for AT only): **NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT M.B.H. [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).**
- (71) Applicant (for all designated States except US): **THE SCRIPPS RESEARCH INSTITUTE [US/US]; 10550 North Torrey Pines Road, La Jolla, CA 92037 (US).**
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **NEMEROW, Glen, R. [US/US]; 462 Cerro Street, Encinitas, CA 92024 (US). LI, Erguang [CN/US]; 7882 Camino Glorita, San Diego, CA 92122 (US).**
- (74) Agent: **BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).**
- (81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**
- (84) Designated States (regional): **ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).**
- Published:**
- without international search report and to be republished upon receipt of that report
  - with sequence listing part of description published separately in electronic form and available upon request from the International Bureau
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 02/04522 A2

(54) Title: **BIFUNCTIONAL MOLECULES AND VECTORS COMPLEXED THEREWITH FOR TARGETED GENE DELIVERY**

(57) Abstract: Methods and products for targeting delivery vectors, such as adenoviral gene delivery particles, to selected cell types are provided. The methods rely on targeting by a bifunctional molecule that specifically complexes with a protein on the vector particle surface and with targeted cell surface proteins. The targeted cell surface proteins are any that activate the phosphatidylinositol-3-OH kinases. The bifunctional molecules, compositions, kits, and methods of preparation and use of the vector/bifunctional molecules for gene therapy are provided.

-1-

## BIFUNCTIONAL MOLECULES AND VECTORS COMPLEXED THEREWITH FOR TARGETED GENE DELIVERY

Work described herein was supported by NIH grants EY11431 and HL54352. The government has certain rights in such subject matter.

### 5 FIELD OF INVENTION

The present invention relates to gene therapy, especially to adenovirus vector-based gene therapy. In particular, adenovirus vectors complexed with bifunctional molecules for targeted delivery of therapeutic and other products are provided. The bifunctional molecules complexed with adenovirus delivery  
10 particles circumvent Coxsackie Adenovirus Receptor (CAR) and integrin interactions and improve gene delivery by such particles. The bifunctional molecules, compositions, kits, and methods of preparation and use of the vector/bifunctional molecules for gene therapy are provided.

### BACKGROUND OF THE INVENTION

#### 15 Adenovirus delivery vectors

Adenovirus, which is a DNA virus with a 36 kilobase (kb) genome, is very well-characterized and its genetics and genetic organization are understood. The genetic organization of adenoviruses permits substitution of large fragments of viral DNA with foreign DNA. In addition, recombinant adenoviruses are  
20 structurally stable and no rearranged viruses are observed after extensive amplification.

Adenoviruses have been employed as delivery vehicles for introducing desired genes into eukaryotic cells. The adenovirus delivers such genes to eukaryotic cells by binding to cellular receptors followed by internalization. The  
25 adenovirus fiber protein is responsible for binding to cells. The fiber protein has two domains, a rod-like shaft portion and a globular head portion that contains the receptor binding region. The fiber spike is a homotrimer, and there are 12 spikes per virion. Human adenoviruses bind to and infect a broad range of cultured cell lines and primary tissues from different species.

30 The 35,000+ base pair (bp) genome of adenovirus type 2 has been sequenced and the predicted amino acid sequences of the major coat proteins (hexon, fiber and penton base) have been described (see, *e.g.*, Neumann *et al.*,

-2-

*Gene* 69: 153-157 (1988); Herisse *et al.*, *Nuc. Acids Res.* 9: 4023-4041 (1981); Roberts *et al.*, *J. Biol. Chem.* 259: 13968-13975 (1984); Kinloch *et al.*, *J. Biol. Chem.* 259: 6431-6436 (1984); and Chroboczek *et al.*, *Viol.* 161: 549-554, 1987).

5           The 35,935 bp sequence of Ad5 DNA is also known and portions of many other adenovirus genomes have been sequenced. The upper packaging limit for adenovirus virions is about 105% of the wild-type genome length (see, e.g., Bett, *et al.*, *J. Virol.* 67(10): 5911-21, 1993). Thus, for Ad2 and Ad5, this would be an upper packaging limit of about 38 kb of DNA.

10           Adenovirus DNA also includes inverted terminal repeat sequences (ITRs) ranging in size from about 100 to 150 bp, depending on the serotype. The inverted repeats permit single strands of viral DNA to circularize by base-pairing of their terminal sequences to form base-paired "panhandle" structures that are required for replication of the viral DNA.

15           For efficient packaging, the ITRs and the packaging signal (a few hundred bp in length) comprise the "minimum requirement" for replication and packaging of a genomic nucleic acid into an adenovirus particle. Helper-dependent vectors lacking all viral ORFs but including these essential *cis* elements (the ITRs and contiguous packaging sequence) have been constructed,

20           Ad vectors have several distinct advantages as gene delivery vehicles. For example, recombination of such vectors is rare; there are no known associations of human malignancies with adenoviral infections despite common human infection with adenoviruses; the genome may be manipulated to accommodate foreign genes of a fairly substantial size; and host proliferation is not required for expression of adenoviral proteins. Adenovirus (Ad)-based gene  
25           delivery vectors efficiently infect many different cells and tissues. This broad tropism, however, means that gene delivery cannot be directed to a specific target cell. A large fraction of intravenously administered adenovirus is retained by the liver, which could lead to undesirable side-effects. Adenovirus may  
30           potentiate immune responses. For example, Adenovirus type 5 (Ad5) also transduces dendritic cells, which present antigens very efficiently, thereby possibly exacerbating the immune response against the vector. It has been

-3-

proposed that vectors with different targeting efficiencies might eliminate these problems, permitting a lower total particle dose and more specific targeting (see, e.g., U.S. application Serial No. 09/482,682).

The wealth of information on adenovirus structure and mechanism of infection, its efficient infection of nondividing cells, and its large genetic capacity make adenovirus a popular gene therapy vector. The wide expression of receptors to which adenovirus binds makes targeting adenovirus vectors difficult. In particular, because of the widespread distribution of the Ad receptor (CAR), current adenoviral (Ad) vectors cannot be targeted to specific cell types (see, e.g., Bergelson *et al.* (1997) *Science* 275:1320-1323; Tomko *et al.* (1997) *Proc.Natl.Acad.Sci. USA* 94:3352-3356). Moreover, CAR and/or internalization receptors ( $\alpha$ v integrins) (see, Wickham *et al.* (1993) *Cell* 73:309-319) are absent or present at low levels on some cell types, rendering them resistant to Ad-mediated gene delivery.

Approximately 20% of all gene therapy clinical trials, registered with the NIH Recombinant DNA Advisory Committee, use replication-deficient adenovirus vectors (Office of Recombinant DNA Database). While successes have been reported, especially in the area of tumor management (see, e.g., Bilbao *et al.* (1998) *Adv.Exp.Med.Biol.* 451:365-374; Gomez-Navarro *et al.* (1999) *Eur.J.Cancer* 35:867-885), the use of adenovirus gene delivery vectors has been hampered by host inflammatory responses to the virus or encoded transgene products (Kay *et al.* (1997) *Proc.Natl.Acad.Sci.USA* 94:12744-12746). In addition, some cell types lack either the Ad attachment receptor, Coxsackie Adenovirus Receptor (CAR; Bergelson *et al.* (1997) *Science* 275:1320-1323; Tomko *et al.* (1997) *Proc.Natl.Acad.Sci. USA* 94:3352-3356) or integrins  $\alpha$ v $\beta$ 3 and  $\alpha$ v $\beta$ 5, which serve as virus internalization receptors (Wickham *et al.* (1993) *Cell* 73:309-319). For example, because airway epithelia do not express CAR and integrins on their apical surface (Goldman *et al.* (1995) *J.Virol.* 69:5951-5958; Grubb *et al.* (1994) *Nature* 371:802-806), clinical trials for the treatment of cystic fibrosis reported variable, generally low, efficacy (Zabner *et al.* (1993) *Cell* 75:207-216; Crystal *et al.* (1994) *Nature genet.* 4:42-51) using adenoviral vectors.



-4-

Thus, while adenoviral vectors and others, hold much promise for therapeutic applications, their usefulness is limited by the widespread tissue distribution of CAR, which restricts delivery to specific cell types, and also by the absence of CAR and/or  $\alpha v$  integrin receptors on certain cells *in vivo*. There have been attempts to overcome these limitations by modifying one or more of the Ad outer capsid proteins in order to retarget vectors to different cell receptor. While improvements in gene delivery have been realized, each method has its limitation. Also, the underlying factors that promote gene delivery have not been clearly defined, which has impeded further progress in Ad vector development; and the specificity of targeting requires further improvement.

Hence, there is a need to improve delivery and targeting of gene delivery vectors, including adenoviral gene delivery vectors, and to understand the underlying basis therefor. Therefore, it is an object herein to provide methods and vectors that can specifically target specific cells and tissues and that provide improved delivery and internalization of such vectors. It is also an object herein to identify the underlying mechanism for internalization and to provide delivery methods and delivery vectors based thereon.

#### SUMMARY OF THE INVENTION

A vector targeting method that takes advantage of the common cell signaling pathways initiated by ligation of  $\alpha v$  integrins and cell surface proteins and receptors that upon ligand interaction activate phosphatidylinositol 3 kinases (PI3K) or the phosphatidylinositol 3 (PI3) signalling pathway is provided. Bifunctional molecules (or multifunctional molecules) for effecting the targeting and complexes that contain the bifunctional molecules conjugated to a viral particle or bacterial surface protein are provided. Methods of gene delivery and gene therapy are also provided. In a preferred embodiment the virus for delivery and to which the bifunctional molecule specifically binds is adenovirus or adeno-associated virus. More preferably the virus is adenovirus, including fiberless viral particles.

In particular, bifunctional molecules that contain an antibody or antigen-binding portion thereof and a targeting agent are provided. The

-5-

antibody specifically binds to an antigen in a protein that binds to  $\alpha_v$  integrin; and the targeting agent specifically binds to a cell surface protein that activates the phosphatidylinositol 3 (PI3) signaling pathway. In particular, the targeting agent binds to a cell surface protein that triggers phosphatidylinositol-3-OH kinase

5 (PI3K) activation.

Thus, the bifunctional molecules include a targeting agent, which is a moiety that specifically binds to a cell surface protein that triggers activation of PI3K, and a binding portion (designated "P" herein) that specifically binds to a protein on a viral particle or bacterial cell surface. Generally such viral or  
 10 bacterial surface protein specifically binds to an  $\alpha_v$  integrin or other protein on the surface of a targeted cell that facilitates or effects internalization of the virus or bacterial into the cell.

The bifunctional molecules optionally include a linker or plurality of linkers that links the antibody or antigen-binding portion thereof to the targeting agent.  
 15 The linker can be a peptide, preferably of from 2 to 100, more preferably 3 to 50, more preferably 5 to 20 amino acids, a single amino acid, or a chemical linker, such as those produced by reaction with crosslinking agents or heterobifunctional crosslinking reagents. The bifunctional molecules can be fusion proteins, chemical conjugates or mixtures thereof, a single amino acid or a  
 20 peptide.

Thus the bifunctional agents can be represented by the formula:

$(P)_n-(L)_q-(TA)_m$ , where  $m$  and  $n$  are integers of 1 or higher, and are generally 1, and  $q$  is 0 or an integer of one or more, and is generally 1 or 2. In  
 25 instances where either or both of  $n$  and  $m$  are greater than 1, the resulting molecule is technically a multifunctional molecule. Each  $P$  and each  $TA$  do can be the same or different.

The protein that binds to  $\alpha_v$  integrin is a viral protein or a bacterial protein that interacts with  $\alpha_v$  integrins for internalization of the respective virus or bacteria. It is with such protein that the "P" moiety of the bifunctional molecule  
 30 interacts. The  $P$  moiety is generally an antibody or portion thereof or recombinant molecule having the binding specificity of an antibody. The antibody or antigen-binding portion of the bifunctional molecule specifically binds

-6-

to such protein, which for example is a viral protein, such as the penton protein of adenovirus.

The antibody or antigen-binding portion of the bifunctional molecule can include a heavy chain or a portion thereof sufficient for antigen-binding fused to the targeting agent; or is an Fab or Fab'2 fragment, or is a synthetic or recombinant molecule that contains antibody fragments, such as portions of the heavy chain and light chain variable regions sufficient for specific interaction with the antigen.

The targeted cell surface protein is any cell surface protein in which binding thereto facilitates internalization, particularly via the signaling pathway that involves PI3K. Such surface proteins, include, but are not limited to cytokine receptors, growth hormone receptors and other non-steroidal hormone receptors, such as a PDGF receptor, an IGF-1 receptor, an EGF receptor, a member of the FGF receptor family, a TNF receptor, a CSF-1 receptor, an insulin receptor, an IGF-1 receptor, an NGF receptor, an IL-2 receptor, an IL-3 receptor, an IL-4 receptor, an IgM receptor, a CD4 receptor, a CD2 receptor, a CD3/T cell receptor, a G protein linked thrombin receptor, an ATP receptor, an fMLP receptor, and tyrosine kinase receptors that, when activated, result in increased accumulation of PtdIns(3,4,5)P<sub>3</sub>, receptors associated with the src family non-receptor tyrosine kinases that stimulate PI3Ks to lead to phosphatidylinositol(3,4,5)P<sub>3</sub> (PtdIns(3,4,5)P<sub>3</sub>) accumulation.

In exemplified embodiments, growth factor/cytokines, such as TNF- $\alpha$ , IGF-1, SCF, PDGF and EGF, hormones, such as insulin, and other molecules or portions thereof that trigger phosphatidylinositol-3-OH kinase (PI3K) activation, a signaling molecule involved in adenovirus internalization, are fused to a monoclonal antibody specific for the viral particle protein that interacts with  $\alpha$ v integrins, which in the case of adenovirus the viral penton base. Ad vectors complexed with these bifunctional mAbs exhibit increased gene delivery of about 10-50 fold to human melanoma cells lacking  $\alpha$ v integrins. The bifunctional Mabs also enhance gene delivery by fiberless adenovirus particles, which cannot bind to CAR. Improved gene delivery correlates with increased virus internalization and attachment as well as PI3K activity. The use of bifunctional mAbs to trigger

-7-

specific cell signaling pathways offers a widely applicable method for bypassing the normal Ad receptors in gene delivery and potentially increasing the selectivity of gene transfer.

Isolated nucleic acid molecule(s) that encode the bifunctional molecules are also provided. In some instances, in which the "P" moiety is an antibody or portion thereof, the bifunctional molecule includes two chains, which can be separately expressed and the reconstituted, such as by the selected expression system. Preferred expression systems are baculovirus systems.

Also provided are combinations that include a bifunctional molecule and a viral or bacterial vector. The components of the combinations may be separate or combined in a single composition in which the bifunctional molecules have been complexed with the viral particles or bacterial cells. Kits, containing the combinations optionally include instructions for administration and/or complexing, are also provided. Upon complexing of a bifunctional molecule and a viral or bacterial vector, the resulting complex can be used for delivery of gene products, such as for therapeutics, gene therapy and or for production of transgenic animals.

Preferred delivery vectors herein are adenovirus vectors, including fiberless adenovirus vectors.

Methods of gene therapy by administration of the targeted gene delivery vectors that include the bifunctional molecules complexed with a viral particle or bacterial cells are also provided.

In exemplified embodiments, bifunctional molecules and complexes thereof with adenovirus delivery vectors are provided. In an exemplary embodiment, a bifunctional molecule that recognizes the RGD motif in the penton base protein of adenovirus is fused to the mature form of a receptor targeting molecule, such as TNF- $\alpha$ . The bifunctional molecules were expressed in insect cells using a non-lytic baculovirus expression system. In particular, the bifunctional molecule was produced from a monoclonal antibody, designated DAV-1, that specifically interacts with penton base on the surface of adenovirus strains, including Ad2, Ad3, Ad4 and Ad5. In particular, this antibody includes sequence of amino acids set forth in SEQ ID No. 2 or SEQ ID No. 6 or a

-8-

sufficient portion thereof for antigen recognition or is encoded by nucleic acid that hybridizes along its full length under conditions of low stringency, more preferably moderate stringency, most preferably high stringency to a sufficient portion of SEQ ID Nos. 1 or 5 to encode an antigen-binding portion of the  
5 antibody. The antibody or portion thereof also include light chain that includes all or a portion of the sequence of amino acids set forth in SEQ ID No. 4 or nucleic acid that hybridizes along its full length under conditions of low stringency, more preferably moderate stringency, most preferably high stringency to a sufficient portion of the SEQ ID No. 3 so that the resulting  
10 molecule encodes an antigen-binding portion of the antibody.

The targeting agent or portion thereof can be selected from among, for example, is a tumor necrosis factor (TNF), a fibroblast growth factor (FGF), an insulin-like growth factor (IGF), a colony stimulating factor (CSF), insulin and a stem cell factor (SCF).

15 In exemplified embodiments, bifunctional molecules were capable of enhancing infection of M21-L12 melanoma cells, which lack  $\alpha v$  integrins and are also resistant to TNF $\alpha$  killing. M21 cells are relatively resistant to transduction with adenovirus vectors. Incubation of Ad encoding lacZ with the original monoclonal antibody alone had little effect enhancing gene delivery. In  
20 contrast, preincubation of Ad.lacZ particles with the bifunctional antibody produced a 20-fold increase in virus-mediated gene delivery. Enhanced virus infection by the bifunctional antibodies was due to a combination of increased virus binding and internalization. Enhanced internalization resulted from increased activation of PI3K through TNF $\alpha$  receptor ligation. Other bifunctional  
25 molecules containing receptor ligands were capable of activating PI3K and enhancing gene delivery.

-9-

These results demonstrate that activation of receptors that activate PI3K bypasses the requirement for  $\alpha v$  integrins or, for adenovirus, CAR to promote virus entry. Receptor bypass was highly effective when cytokine or growth factors were activated in close proximity to bound virus particles.

## 5 DETAILED DESCRIPTION OF THE INVENTION

### A. DEFINITIONS

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which this invention belongs. All patents, applications, published applications  
10 and other publications and sequences from GenBank and other data bases referred to anywhere in the disclosure herein are incorporated by reference in their entirety.

As used herein, the amino acids, which occur in the various amino acid sequences appearing herein, are identified according to their well-  
15 known, three-letter or one-letter abbreviations. The nucleotides, which occur in the various DNA fragments, are designated with the standard single-letter designations used routinely in the art (see, Table 1).

As used herein, amino acid residue refers to an amino acid formed upon chemical digestion (hydrolysis) of a polypeptide at its peptide linkages. The  
20 amino acid residues described herein are preferably in the "L" isomeric form. However, residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the desired functional property is retained by the polypeptide.  $\text{NH}_2$  refers to the free amino group present at the amino terminus of a polypeptide.  $\text{COOH}$  refers to the free carboxy group present at the carboxyl  
25 terminus of a polypeptide. In keeping with standard polypeptide nomenclature described in *J. Biol. Chem.*, 243:3552-59 (1969) and adopted at 37 C.F.R. § § 1.821 - 1.822, abbreviations for amino acid residues are shown in the following Table:

-10-

Table 1  
Table of Correspondence

SYMBOL		
1-Letter	3-Letter	AMINO ACID
Y	Tyr	tyrosine
G	Gly	glycine
F	Phe	phenylalanine
M	Met	methionine
A	Ala	alanine
S	Ser	serine
I	Ile	isoleucine
L	Leu	leucine
T	Thr	threonine
V	Val	valine
P	Pro	proline
K	Lys	lysine
H	His	histidine
Q	Gln	glutamine
E	Glu	glutamic acid
Z	Glx	Glu and/or Gln
W	Trp	tryptophan
R	Arg	arginine
D	Asp	aspartic acid
N	Asn	asparagine
B	Asx	Asn and/or Asp
C	Cys	cysteine
X	Xaa	Unknown or other

It should be noted that all amino acid residue sequences represented  
herein by formulae have a left to right orientation in the conventional direction of

-11-

amino-terminus to carboxyl-terminus. In addition, the phrase "amino acid residue" is broadly defined to include the amino acids listed in the Table of Correspondence and modified and unusual amino acids, such as those referred to in 37 C.F.R. § § 1.821-1.822, and incorporated herein by reference.

- 5 Furthermore, it should be noted that a dash at the beginning or end of an amino acid residue sequence indicates a peptide bond to a further sequence of one or more amino acid residues or to an amino-terminal group such as NH<sub>2</sub> or to a carboxyl-terminal group such as COOH.

- In a peptide or protein, suitable conservative substitutions of amino acids are known to those of skill in this art and may be made generally without altering the biological activity of the resulting molecule. Those of skill in this art recognize that, in general, single amino acid substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, e.g., Watson et al. *Molecular Biology of the Gene*, 4th Edition, 1987, The Bejacmin/Cummings Pub. co., p.224).

Such substitutions are preferably made in accordance with those set forth in TABLE 2 as follows:

TABLE 2		
	Original residue	Conservative substitution
20	Ala (A)	Gly; Ser
	Arg (R)	Lys
	Asn (N)	Gln; His
	Cys (C)	Ser
	Gln (Q)	Asn
25	Glu (E)	Asp
	Gly (G)	Ala; Pro
	His (H)	Asn; Gln
	Ile (I)	Leu; Val
	Leu (L)	Ile; Val
30	Lys (K)	Arg; Gln; Glu
	Met (M)	Leu; Tyr; Ile
	Phe (F)	Met; Leu; Tyr
	Ser (S)	Thr
	Thr (T)	Ser
35	Trp (W)	Tyr
	Tyr (Y)	Trp; Phe
	Val (V)	Ile; Leu

Other substitutions are also permissible and may be determined empirically or in accord with known conservative substitutions.



-12-

As used herein, a complementing plasmid describes plasmid vectors that deliver nucleic acids into a packaging cell line for stable integration into a chromosome in the cellular genome.

As used herein, a delivery plasmid is a plasmid vector that carries or  
5 delivers nucleic acids encoding a therapeutic gene or gene that encodes a therapeutic product or a precursor thereof or a regulatory gene or other factor that results in a therapeutic effect when delivered *in vivo* in or into a cell line, such as, but not limited to a packaging cell line, to propagate therapeutic viral vectors.

10 As used herein, a variety of vectors are described. For example, one vector is used to deliver particular nucleic acid molecules into a packaging cell line for stable integration into a chromosome. These types of vectors are generally identified herein as complementing plasmids. A further type of vector described herein carries or delivers nucleic acid molecules in or into a cell line  
15 (e.g., a packaging cell line) for the purpose of propagating therapeutic viral vectors; hence, these vectors are generally referred to herein as delivery plasmids. A third "type" of vector described herein is used to carry nucleic acid molecules encoding therapeutic proteins or polypeptides or regulatory proteins or are regulatory sequences to specific cells or cell types in a subject in need of  
20 treatment; these vectors are generally identified herein as therapeutic viral vectors or recombinant adenoviral vectors or viral Ad-derived vectors and are in the form of a virus particle encapsulating a viral nucleic acid containing an expression cassette for expressing the therapeutic gene.

As used herein, a DNA or nucleic acid homolog refers to a nucleic acid  
25 that includes a preselected conserved nucleotide sequence, such as a sequence encoding a therapeutic polypeptide. By the term "substantially homologous" is meant having at least 80%, preferably at least 90%, most preferably at least 95% homology therewith or a less percentage of homology or identity and conserved biological activity or function.

30 The terms "homology" and "identity" are often used interchangeably. In this regard, percent homology or identity may be determined, for example, by comparing sequence information using a GAP computer program. The GAP

-13-

program utilizes the alignment method of Needleman and Wunsch (*J. Mol. Biol.* 48:443 (1970)), as revised by Smith and Waterman (*Adv. Appl. Math.* 2:482 (1981)). Briefly, the GAP program defines similarity as the number of aligned symbols (i.e., nucleotides or amino acids) which are similar, divided by the total number of symbols in the shorter of the two sequences. The preferred default parameters for the GAP program may include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) and the weighted comparison matrix of Gribskov and Burgess, *Nucl. Acids Res.* 14:6745 (1986), as described by Schwartz and Dayhoff, eds., *ATLAS OF PROTEIN SEQUENCE AND STRUCTURE*, National Biomedical Research Foundation, pp. 353-358 (1979); (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps.

Whether any two nucleic acid molecules have nucleotide sequences that are at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% "identical" can be determined using known computer algorithms such as the "FAST A" program, using for example, the default parameters as in Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444 (1988). Alternatively the BLAST function of the National Center for Biotechnology Information database may be used to determine identity

In general, sequences are aligned so that the highest order match is obtained. "Identity" *per se* has an art-recognized meaning and can be calculated using published techniques. (See, e.g.: *Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part I*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991).

While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans (Carillo, H. & Lipton, D., *SIAM J Applied Math* 48:1073 (1988)).

-14-

Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H. & Lipton, D., *SIAM J Applied Math* 48:1073 (1988). Methods to  
5 determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(11):387 (1984)), BLASTP, BLASTN, FASTA (Atschul, S.F., et al., *J Molec Biol* 215:403 (1990)).

10 Therefore, as used herein, the term "identity" represents a comparison between a test and a reference polypeptide or polynucleotide. For example, a test polypeptide may be defined as any polypeptide that is 90% or more identical to a reference polypeptide.

As used herein, the term at least "90% identical to" refers to percent  
15 identities from 90 to 99.99 relative to the reference polypeptides. Identity at a level of 90% or more is indicative of the fact that, assuming for exemplification purposes a test and reference polynucleotide length of 100 amino acids are compared. No more than 10% (i.e., 10 out of 100) amino acids in the test polypeptide differs from that of the reference polypeptides. Similar comparisons  
20 may be made between a test and reference polynucleotides. Such differences may be represented as point mutations randomly distributed over the entire length of an amino acid sequence or they may be clustered in one or more locations of varying length up to the maximum allowable, e.g. 10/100 amino acid difference (approximately 90% identity). Differences are defined as nucleic  
25 acid or amino acid substitutions, or deletions.

As used herein, stringency conditions refer to the washing conditions for removing the non-specific probes and conditions that are equivalent to either high, medium, or low stringency as described below:

- 1) high stringency: 0.1 x SSPE, 0.1% SDS, 65°C
- 30 2) medium stringency: 0.2 x SSPE, 0.1% SDS, 50°C
- 3) low stringency: 1.0 x SSPE, 0.1% SDS, 50°C.

-15-

It is understood that equivalent stringencies may be achieved using alternative buffers, salts and temperatures.

As used herein, genetic therapy involves the transfer of heterologous DNA to the certain cells, target cells, of a mammal, particularly a human, with a disorder or conditions for which such therapy is sought. The DNA is introduced into the selected target cells in a manner such that the heterologous DNA is expressed and a therapeutic product encoded thereby is produced.

Alternatively, the heterologous DNA may in some manner mediate expression of DNA that encodes the therapeutic product, it may encode a product, such as a peptide or RNA that in some manner mediates, directly or indirectly, expression of a therapeutic product. Genetic therapy may also be used to nucleic acid encoding a gene product replace a defective gene or supplement a gene product produced by the mammal or the cell in which it is introduced. The introduced nucleic acid may encode a therapeutic compound, such as a growth factor inhibitor thereof, or a tumor necrosis factor or inhibitor thereof, such as a receptor therefor, that is not normally produced in the mammalian host or that is not produced in therapeutically effective amounts or at a therapeutically useful time. The heterologous DNA encoding the therapeutic product may be modified prior to introduction into the cells of the afflicted host in order to enhance or otherwise alter the product or expression thereof.

As used herein, heterologous DNA is DNA that encodes RNA and proteins that are not normally produced *in vivo* by the cell in which it is expressed or that mediates or encodes mediators that alter expression of endogenous DNA by affecting transcription, translation, or other regulatable biochemical processes.

Heterologous DNA may also be referred to as foreign DNA. Any DNA that one of skill in the art would recognize or consider as heterologous or foreign to the cell in which is expressed is herein encompassed by heterologous DNA.

Examples of heterologous DNA include, but are not limited to, DNA that encodes traceable marker proteins, such as a protein that confers drug resistance, DNA that encodes therapeutically effective substances, such as anti-cancer agents, enzymes and hormones, and DNA that encodes other types of proteins, such as antibodies. Antibodies that are encoded by heterologous DNA may be secreted

-16-

or expressed on the surface of the cell in which the heterologous DNA has been introduced.

Hence, herein heterologous DNA or foreign DNA, refers to a DNA molecule not present in the exact orientation and position as the counterpart  
5 DNA molecule found in the corresponding wild-type adenovirus. It may also refer to a DNA molecule from another organism or species (*i.e.*, exogenous) or from another Ad serotype.

As used herein, a therapeutically effective product is a product that is encoded by heterologous DNA that, upon introduction of the DNA into a host, a  
10 product is expressed that effectively ameliorates or eliminates the symptoms, manifestations of an inherited or acquired disease or that cures said disease.

Typically, DNA encoding the desired heterologous DNA is cloned into a plasmid vector and introduced by routine methods, such as calcium-phosphate mediated DNA uptake (see, (1981) Somat. Cell. Mol. Genet. 7:603-616) or  
15 microinjection, into producer cells, such as packaging cells. After amplification in producer cells, the vectors that contain the heterologous DNA are introduced into selected target cells.

As used herein, an expression or delivery vector refers to any plasmid or virus into which a foreign or heterologous DNA may be inserted for expression in  
20 a suitable host cell — *i.e.*, the protein or polypeptide encoded by the DNA is synthesized in the host cell's system. Vectors capable of directing the expression of DNA segments (genes) encoding one or more proteins are referred to herein as "expression vectors." Also included are vectors that allow cloning of cDNA (complementary DNA) from mRNA produced using reverse  
25 transcriptase.

As used herein, a gene is a nucleic acid molecule whose nucleotide sequence encodes RNA or polypeptide. A gene can be either RNA or DNA. Genes may include regions preceding and following the coding region (leader and trailer) as well as intervening sequences (introns) between individual coding  
30 segments (exons).

-17-

As used herein, tropism with reference to a adenovirus refers to the selective infectivity or binding that is conferred on the particle by the fiber protein, particularly the C-terminus portion that comprises the knob.

As used herein, isolated with reference to a nucleic acid molecule or polypeptide or other biomolecule means that the nucleic acid or polypeptide has separated from the genetic environment from which the polypeptide or nucleic acid were obtained. It may also mean altered from the natural state. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Thus, a polypeptide or polynucleotide produced and/or contained within a recombinant host cell is considered isolated. Also intended as an "isolated polypeptide" or an "isolated polynucleotide" are polypeptides or polynucleotides that have been purified, partially or substantially, from a recombinant host cell or from a native source. For example, a recombinantly produced version of a compounds can be substantially purified by the one-step method described in Smith and Johnson, *Gene* 67:31-40 (1988). The terms isolated and purified are sometimes used interchangeably.

Thus, by "isolated" is meant that the nucleic is free of the coding sequences of those genes that, in the naturally-occurring genome of the organism (if any) immediately flank the gene encoding the nucleic acid of interest. Isolated DNA may be single-stranded or double-stranded, and may be genomic DNA, cDNA, recombinant hybrid DNA, or synthetic DNA. It may be identical to a native DNA sequence, or may differ from such sequence by the deletion, addition, or substitution of one or more nucleotides.

Isolated or purified as it refers to preparations made from biological cells or hosts means any cell extract containing the indicated DNA or protein including a crude extract of the DNA or protein of interest. For example, in the case of a protein, a purified preparation can be obtained following an individual technique or a series of preparative or biochemical techniques and the DNA or protein of interest can be present at various degrees of purity in these preparations. The procedures may include for example, but are not limited to, ammonium sulfate

-18-

fractionation, gel filtration, ion exchange change chromatography, affinity chromatography, density gradient centrifugation and electrophoresis.

A preparation of DNA or protein that is "substantially pure" or "isolated" should be understood to mean a preparation free from naturally occurring materials with which such DNA or protein is normally associated in nature. "Essentially pure" should be understood to mean a "highly" purified preparation that contains at least 95% of the DNA or protein of interest.

A cell extract that contains the DNA or protein of interest should be understood to mean a homogenate preparation or cell-free preparation obtained from cells that express the protein or contain the DNA of interest. The term "cell extract" is intended to include culture media, especially spent culture media from which the cells have been removed.

As used herein, a packaging cell line is a cell line that provides a missing gene product or its equivalent.

As used herein, an adenovirus viral particle is the minimal structural or functional unit of a virus. A virus can refer to a single particle, a stock of particles or a viral genome. The adenovirus (Ad) particle is relatively complex and may be resolved into various substructures.

As used herein, "penton" or "penton complex" are preferentially used herein to designate a complex of penton base and fiber. The term "penton" may also be used to indicate penton base, as well as penton complex. The meaning of the term "penton" alone should be clear from the context within which it is used.

As used herein, a plasmid refers to an autonomous self-replicating extrachromosomal circular nucleic acid molecule, typically DNA.

As used herein, a post-transcription regulatory element (PRE) is a regulatory element found in viral or cellular messenger RNA that is not spliced, i.e. intronless messages. Examples include, but are not limited to, human hepatitis virus, woodchuck hepatitis virus, the TK gene and mouse histone gene. The PRE may be placed before a polyA sequence and after a heterologous DNA sequence.

-19-

As used herein, pseudotyping describes the production of adenoviral vectors having modified capsid protein or capsid proteins from a different serotype than the serotype of the vector itself. One example, is the production of an adenovirus 5 vector particle containing an Ad37 fiber protein. This may be accomplished by producing the adenoviral vector in packaging cell lines expressing different fiber proteins.

As used herein, promoters of interest herein may be inducible or constitutive. Inducible promoters will initiate transcription only in the presence of an additional molecule; constitutive promoters do not require the presence of any additional molecule to regulate gene expression. a regulatable or inducible promoter may also be described as a promoter where the rate or extent of RNA polymerase binding and initiation is modulated by external stimuli. Such stimuli include, but are not limited to various compounds or compositions, light, heat, stress and chemical energy sources. Inducible, suppressible and repressible promoters are considered regulatable promoters. Preferred promoters herein, are promoters that are selectively expressed in ocular cells, particularly photoreceptor cells.

As used herein, receptor refers to a biologically active molecule that specifically binds to (or with) other molecules. The term "receptor protein" may be used to more specifically indicate the proteinaceous nature of a specific receptor.

As used herein, recombinant refers to any progeny formed as the result of genetic engineering. This may also be used to describe a virus formed by recombination of plasmids in a packaging cell.

As used herein, a transgene or therapeutic nucleic acid molecule includes DNA and RNA molecules encoding an RNA or polypeptide. Such molecules may be "native" or naturally-derived sequences; they may also be "non-native" or "foreign" that are naturally- or recombinantly-derived. The term "transgene," which may be used interchangeably herein with the term "therapeutic nucleic acid molecule," is often used to describe a heterologous or foreign (exogenous) gene that is carried by a viral vector and transduced into a host cell.



-20-

Therefore, therapeutic nucleotide nucleic acid molecules include antisense sequences or nucleotide sequences which may be transcribed into antisense sequences. Therapeutic nucleotide sequences (or transgenes) all include nucleic acid molecules that function to produce a desired effect in the cell or cell nucleus  
5 into which said therapeutic sequences are delivered. For example, a therapeutic nucleic acid molecule can include a sequence of nucleotides that encodes a functional protein intended for delivery into a cell which is unable to produce that functional protein.

As used herein, a promoter region refers to the portion of DNA of a gene  
10 that controls transcription of the DNA to which it is operatively linked. The promoter region includes specific sequences of DNA that are sufficient for RNA polymerase recognition, binding and transcription initiation. This portion of the promoter region is referred to as the promoter. In addition, the promoter region includes sequences that modulate this recognition, binding and transcription  
15 initiation activity of the RNA polymerase. These sequences may be *cis* acting or may be responsive to *trans* acting factors. Promoters, depending upon the nature of the regulation, may be constitutive or regulated.

As used herein, the phrase "operatively linked" generally means the sequences or segments have been covalently joined into one piece of DNA,  
20 whether in single or double stranded form, whereby control sequences on one segment control expression or replication or other such control of other segments. The two segments are not necessarily contiguous.

As used herein, exogenous encompasses any therapeutic composition that is administered by the therapeutic methods provided herein. Thus,  
25 exogenous may also be referred to herein as foreign, or non-native or other equivalent expression.

As used herein, antibody refers to an immunoglobulin, whether natural or partially or wholly synthetically produced, including any derivative thereof that retains the specific binding ability the antibody. Hence antibody includes any  
30 protein having a binding domain that is homologous or substantially homologous to an immunoglobulin binding domain. Antibodies include members of any immunoglobulin claims, including IgG, IgM, IgA, IgD and IgE.

-21-

As used herein, antibody fragment refers to any derivative of an antibody that is less than full length, retaining at least a portion of the full-length antibody's specific binding ability. Examples of antibody fragments include, but are not limited to, Fab, Fab', F(ab)<sub>2</sub>, single-chain Fvs (scFV), FV, dsFV diabody and Fd fragments and other recombinant form of antibodies that retain or exhibit binding specificity.

The fragment can include multiple chains linked together, such as by disulfide bridges. An antibody fragment generally contains at least about 50 amino acids and typically at least 200 amino acids. For purposes herein, any fragment that retains specific binding for a  $\alpha$ v integrin binding protein is contemplated. Such fragments may be produced by chemical or recombinant means.

As used herein, an Fv antibody fragment is composed of one variable heavy domain (V<sub>H</sub>) and one variable light domain linked by noncovalent interactions.

As used herein, a dsFV refers to an Fv with an engineered intermolecular disulfide bond, which stabilizes the V<sub>H</sub>-V<sub>L</sub> pair.

As used herein, an F(ab)<sub>2</sub> fragment is an antibody fragment that results from digestion of an immunoglobulin with pepsin at pH 4.0-4.5; it may be recombinantly produced.

Thus, with reference to the DAV-1 antibody exemplified herein, the Fab portion is the entire light chain and amino acids 1-229 of the DAV-1 heavy chain (SEQ ID Nos. 1 and 2). The Fab fragment is involved in antigen binding, with the exception of the first 19 amino acids which constitute the signal peptide sequence for secretion of the antibody. Fab fragments can be generated by digestion with papain.

The Fab'2 portion is the Fab fragment and the hinge region, which connects the Fab antigen-binding portion with the Fc portion which is involved in complement activation and macrophage binding. Amino acids 230-242 of the DAV-1 heavy fragment constitute the hinge region of DAV-1. The Fab'2 fragments used in this manuscript that were generated by cloning comprise

-22-

amino acids 1-247 of the DAV-1 heavy chain sequence. Fab'2 fragments can be generated by digestion with pepsin.

As used herein, Fab fragments is an antibody fragment that results from digestion of an immunoglobulin with papain; it may be recombinantly produced.

5 As used herein, scFVs refer to antibody fragments that contain a variable light chain ( $V_L$ ) and variable heavy chain ( $V_H$ ) covalently connected by a polypeptide linker in any order. The linker is of a length such that the two variable domains are bridged without substantial interference. Preferred linkers are  $(\text{Gly-Ser})_n$  residues with some Glu or Lys residues dispersed throughout to  
10 increase solubility.

As used herein, diabodies are dimeric scFV; diabodies typically have shorter peptide linkers than scFVs, and they preferentially dimerize.

As used herein, humanized antibodies refer to antibodies that are modified to include "human" sequences of amino acids so that administration to  
15 a human will not provoke an immune response. Methods for preparation of such antibodies are known. For example, the hybridoma that expresses the monoclonal antibody is altered by recombinant DNA techniques to express an antibody in which the amino acid composition of the non-variable regions is based on human antibodies. Computer programs have been designed to identify  
20 such regions. For human therapy, humanized antibodies are preferred.

As used herein, production by recombinant means by using recombinant DNA methods means the use of the well known methods of molecular biology for expressing proteins encoded by cloned DNA, including cloning expression of genes and methods, such as gene shuffling and phage display with screening for  
25 desired specificities.

As used herein, the term "conjugated" refers stable attachment, such ionic or covalent attachment.

As used herein, a composition refers to any mixture of two or more products or compounds. It may be a solution, a suspension, liquid, powder, a  
30 paste, aqueous, non-aqueous or any combination thereof.

-23-

As used herein, a combination refers to any association between two or more items.

As used herein, fluid refers to any composition that can flow. Fluids thus encompass compositions that are in the form of semi-solids, pastes, solutions, aqueous mixtures, gels, lotions, creams and other such compositions.

As used herein, substantially identical to a product means sufficiently similar so that the property of interest is sufficiently unchanged so that the substantially identical product can be used in place of the product.

#### B. Adenovirus delivery and internalization

##### Attachment and internalization

Adenovirus attachment to cells is mediated by the elongated fiber, which mediates initial attachment of the virus to cell receptors. The fiber is a homotrimeric protein with an elongated central shaft domain that varies in length among different virus serotypes. The C terminus of the protein contains a globular domain known as the knob, which is responsible for receptor interaction. The N terminus anchors the protein to the virus surface via interaction with the penton base protein.

The cellular receptor recognized by fiber is known as Coxsackie Adenovirus Receptor (CAR), which is a member of the Ig superfamily and contains two extracellular domains. The N-terminal domain is sufficient to mediate fiber interaction. Secondary interactions of the virus penton base protein with  $\alpha$ v integrins promote virus internalization into clathrin-coated pits and endosomes.

Virions are capable of disrupting the early endosome allowing the majority of the virion particle to escape into the cytoplasm. Viral particles are then transported along microtubules to the nuclear pore complex and virion DNA is translocated into the nucleus.

Ad enter cells by a clathrin-coated pit pathway. Ad internalization is a complex process that is found herein to be similar in some respects to that described for cell invasion by certain pathogenic bacteria. Clustering of  $\alpha$ v integrins (not CAR) results in the formation of a cytoplasmic signaling complex involving at least three major components (cSRC, CAS, PI3K). This signaling

-24-

complex is capable of further activation of the Rho family of small GTPases (RAC, CDC42) whose activation ultimately results in the reorganization of actin cytoskeleton and enhanced virus internalization.

Adenovirus internalization via the clathrin-coated pit pathway (Patterson  
5 *et al.* (1983) *J.Gen.Virol.* 64:1091-1099; Wang *et al.* (1998) *J.Virol.* 72:3455-  
3458 ) requires activation of several signaling molecules including  
phosphatidylinositol 3-OH kinase (PI3K; Li *et al.* (1998) *J.Virol.* 72:2055-2061)  
and the Rho family of small GTPases (Li *et al.* (1998) *J.Virol.* 72:8806-8812).  
The major downstream target of this signaling pathway is the actin cytoskeleton,  
10 which promotes virus uptake.

#### PI3Ks

The phosphatidylinositol 3-kinases (PI3 kinases or PI3Ks) represent a  
ubiquitous family of heterodimeric lipid kinases that are found in association with  
the cytoplasmic domain of hormone and growth factor receptors and oncogene  
15 products. Phosphoinositide 3-OH-kinases (PI3Ks) constitute a large family of  
enzymes capable of 3-phosphorylating at least one of the cellular  
phosphoinositides (Whitman *et al.* (1988) *Nature* 332:644-646).  
3-phosphorylated phosphoinositides are found in all higher eukaryotic cells.

PI3Ks act as downstream effectors of the above-noted receptors, are  
20 recruited upon receptor stimulation and mediate the activation of second  
messenger signaling pathways through the production of phosphorylated deriva-  
tives of inositol. PI3Ks have been implicated in many cellular activities including  
growth factor mediated cell transformation, mitogenesis, protein trafficking, cell  
survival and proliferation, DNA synthesis, apoptosis, neurite outgrowth, nitric  
25 oxide signaling and insulin-stimulated glucose transport.

PI3Ks phosphorylate phosphatidylinositol (PtdIns) at the 3'-hydroxyl of  
the inositol ring and substrates include PtdIns, PtdIns(4)phosphate PtdIns(4,5)bi-  
phosphate and PtdIns(3,4,5) triphosphate; the major product is PtdIns(3,4,5)tri-  
phosphate (see, *e.g.*, Fry (1994) *Biochim. Biophys. Acta.* 1226: 237-268). A  
30 PI3K and a lipid product of this enzyme, phosphatidylinositol (3,4,5)-triphosphate  
(hereinafter "PtdIns(3,4,5)P3"), are part of an important second messenger

-25-

system in cellular signal transduction. PtdIns(3,4,5) P3 appears to be a second messenger in extremely diverse signalling pathways.

As described herein, PI3K is a member of a family of lipid kinases that include a p85 regulatory subunit and a p110 catalytic subunit (Toker *et al.*,  
5 *Nature*, 387:673 (1997)). The p85 subunit of PI3K binds directly to phosphorylated FAK (Chen *et al.*, *J. Biol. Chem.*, 271:26329 (1996)). The products of PI3K activation, phosphatidylinositol-3,4-bisphosphate and phosphatidylinositol-3,4,5-trisphosphate are increased in the plasma membrane of activated but not quiescent cells and have been proposed to act as second  
10 messengers for a number of cell functions including cell motility, the Ras pathway, vesicle trafficking and secretion, apoptosis, the movement of organelle membranes, shape alteration through rearrangement of cytoskeletal actin, transformation, chemotaxis, cell cycle progression and intracellular protein trafficking (Carpenter *et al.*, *Curr. Opin. Cell Biol.*, 8:153 (1996); Chou *et al.*,  
15 *Cell* 85:573 (1996); Wennström *et al.*, *Curr. Biol.*, 4:385 (1994); Toker *et al.*, *Nature*, 387:673 (1997)). The phospholipid second messengers phosphatidylinositol-3,4-bisphosphate and phosphatidylinositol-3,4,5-trisphosphate mediate the cell functions and processes by activation of protein kinase B and the small GTP-binding proteins Ras, Rho, Rac and CDC42 (Hall, *Science*,  
20 279:509 (1998); Karlund *et al.*, *Science* 275:1927 (1997); Tapon *et al.*, *Curr. Opin. Cell Biol.*, 9:86 (1995)). Activation of Rac and CDC42 induces polymerization of monomeric actin, resulting in the formation of a dense network of actin filaments underlying the plasma membrane, and the actin-rich regions form a variety of membrane extensions known as lamellipodia ("membrane  
25 ruffling") and filopodia ("hairlike structures") (Luo *et al.*, *Nature* 379:837 (1996)). Activation of Rho is associated with the formation of actin-associated stress fibers (Nobes *et al.*, *Cell*, 81:53 (1995); Ridley *et al.*, *Cell*, 70:389 (1992)). The polymerized actin filaments maintain the architecture of the surface protrusions. This actin assembly initiated by activation of PI3K plays an  
30 important role in a number of biological and/or pathogenic processes including directed cell migration (Zigmond, *Curr. Biol.*, 8:66 (1996)), axonal guidance in neuronal cells (Luo *et al.*, *Genes Dev.*, 8:1787 (1994)), and cell invasion by a

-26-

number of different bacteria (Chen *et al.*, *Science*, 274:2115 (1996); Hordijk *et al.*, *Science*, 278:1464 (1998)). Recent studies have indicated that actin polymerization also facilitates clathrin-mediated endocytosis (Benmerah *et al.*, *J. Cell Biol.*, 140:1055 (1998); Lamaze *et al.*, *J. Biol. Chem.*, 272:20332 (1997);

5 Witke *et al.*, *EMBO J.*, 17:967 (1998)).

A number of stimulatory agonists activate the PI3K-mediated signalling pathway. Many of the effects of insulin on glucose and lipid metabolism are elicited through activation of PI3K (Shepherd *et al.*, *Biochem. J.*, 333(3):471 (1998)). IGF-1, PDGF and EGF stimulate cell proliferation in astroglial cells  
10 through increased PI3K activity (Pomerance *et al.*, *J. Neurosci. Res.*, 40:737 (1995)). Insulin or IGF-1 induced membrane ruffling is mediated via activation of PI3K (Kotani *et al.*, *EMBO J.*, 13(10):2313 (1994)). PI3K-associated p85 associates with EGFR and PDGFR upon stimulation with EGF or PDGF, respectively (Hu *et al.*, *Mol. Cell. Biol.*, 12:981 (1992)). Fibronectin, insulin and  
15 PDGF stimulation of ILK (integrin-linked kinase) are dependent on the activity of PI3K (Delcommenne *et al.*, *Proc. Natl. Acad. Sci. USA* 95:11211 (1998)). Stimulation of certain cell types with TNF- $\alpha$  has been shown to activate PI3K (Guo *et al.*, *J. Biol. Chem.*, 271:615 (1996)). PI3K is also activated by internalins, which are bacterial surface proteins involved in bacterial invasion  
20 (Ireton *et al.*, *Science*, 274:780 (1996)).

Adenovirus endocytosis also requires activation of PI3K (Li *et al.*, *J. Virol.* 72:2055 (1998)). Adenovirus endocytosis is a multistep process beginning with its attachment to cells via the elongated fiber protein which the cell surface protein receptor known as CAR (Bergelson *et al.*, *Science*, 275:1320 (1997)).  
25 Secondary interactions of the virus penton base protein via their RGD motifs with  $\alpha_v$  integrins on the cell surface promotes virus internalization, via a receptor-mediated endocytosis pathway, into clathrin-coated pits and endosomes. While fiber binding is not required for PI3K activation, the interaction of adenovirus penton base protein to with  $\alpha_v$  integrins stimulates  
30 PI3K (Li *et al.*, *J. Virol.* 72:2055 (1998)). Stimulation of PI3K has been shown to be essential to viral entry; when cells were treated with the PI3K-specific inhibitors wortmannin and LY294002, adenovirus internalization was inhibited (Li

-27-

*et al.*, *J. Virol.* 72:2055 (1998)). The clustering of the  $\alpha$  integrins results in the formation of a cytoplasmic signaling complex involving at least three major components: cSRC, CAS and PI3K. This signaling complex is capable of further activation of the Rho family of small GTPases (Rac, CDC42) whose activation ultimately results in the reorganization of the actin cytoskeleton and enhanced virus internalization.

Thus, the signaling pathways activated by growth factors/cytokine receptors, including tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ), insulin-like growth factor 1 (IGF-1) and epidermal growth factor (EGF) receptors (Hotamisligil *et al.* (1994) *Proc.Natl.Acad.Sci.USA* 24:4854-4858; Guo *et al.* (1996) *J.Biol.Chem.* 271:615-618; Pomerance *et al.* (1995) *J.Neurosci.* 40:737-746; Kotani, *et al.* (1994) *EMBO J.* 13:2313-2321; and Hu *et al.* (1992) *Mol.Cell Biol.* 12:981-990) initiate signaling events that converge at PI3K activation.

It is shown herein that advantage can be taken of the similarity in signaling processes elicited by cell surface receptors that activate PI3 kinases and the  $\alpha$  integrin mediated viral internalization, providing the basis for the methods and vectors provided herein, which bypass Ad integrin interaction to facilitate gene delivery. It is shown herein that molecules that bind to and activate receptors that participate in the PI3 signalling pathway can be used to target linked moieties, such as adenoviral particles, to such receptors, which include hormone and growth factor receptors, other G-protein coupled receptors and receptors for various oncogenes, and effect internalization.

#### 1. Bifunctional molecules

Provided herein are bifunctional molecules that are conjugates of an agent, such as an antibody or fragment thereof, that specifically binds to  $\alpha$  integrin-binding protein (referred to as component "P" herein) and an agent (referred to as TA) that targets moieties linked to the protein that specifically binds to the  $\alpha$  integrin-binding protein to cells that express surface receptors to which the targeting agent specifically binds. TA and P are joined directly, typically by covalent linkage, or are linked via a linker L.

For purposes herein these conjugates are referred to as bifunctional molecules and include the following components: (TA)<sub>n</sub>, P<sub>m</sub> and L<sub>q</sub>, wherein n



-28-

and m are integers of 1 or more, typically 1 or 2, and q is 0 or an integer of 1 or more. These components are selected such that the resulting molecules will bind to or interact with a protein on a viral particle or bacteria to form a complex and target the resulting complex to a cell surface protein that activates PI3K and that  
5 is recognized by the targeting agent such that the resulting complex is internalized. Where n and m are other than one, the resulting molecules are multifunctional, in such instances more than one TA and/or P moiety can be used, each may be the same or different. The components TA, P and L may be conjugated by covalent linkages, ionic linkages or any other bond resulting in  
10 attachment.

It is understood that the P and the targeting agent (or linker and targeted agent) may be linked in any order and through any appropriate linkage, as long as the resulting conjugate binds to a receptor to which targeted binds and internalizes the complexed viral particles or bacterial cells in cells bearing the  
15 receptor.

Generally, P is an antibody or fragment thereof that specifically binds to  $\alpha v$  integrin binding proteins, such as the penton protein of adenovirus. P is preferably a monoclonal antibody or fragment thereof or synthetic antibody or recombinant protein that includes a sufficient portion of the antibody chains to  
20 specifically bind to a selected antigen. In this instance, the antigen is one present in a protein on a viral particle or bacterial surface that mediates attachment of the viral particle of bacterium to  $\alpha v$  integrins.

a. Antigen-binding portion

In preferred embodiments, the moiety (P) that binds to proteins that  
25 interact with  $\alpha v$  integrins, is a monoclonal antibody or antigen-binding fragment thereof. Proteins, such as the penton protein of adenovirus, that interact with  $\alpha v$  integrins to facilitate internalization of viral particles and bacteria occur on viral particles and bacterial cells surfaces. Antibodies specific therefor can be made by standard methods, such as using hybridoma technology or by making  
30 recombinant antibodies or fragments thereof and screening, such as through phage display technology, for binding to the viral or bacterial surface protein. The antibody or fragment thereof or recombinant version thereof, is then

-29-

modified to include a targeting agent, such as a growth factor or hormone or other protein that binds to receptors that activate the PI3K signaling pathway. The resulting bifunctional molecule specifically binds to a viral or bacterial surface protein, such as penton, and also will specifically bind to a selected  
5 targeted cell that expresses the targeted receptors.

b. Targeting agents

The targeting agents are those that are ligands, including hormones, growth factors and cytokines, that specifically bind to and activate receptors that activate the PI3K signaling pathway. Hence the targeting agent is chosen  
10 to bind to receptors. Selecting targeted receptors can be those that are overexpressed in a particular disorder, such as an angiogenic disorder, including cancers, and inflammatory disorders. PI3K activity in cells of hematopoietic lineage, particularly neutrophils, monocytes, and other types of leukocytes, is involved in many of the non-memory immune responses associated with acute  
15 and chronic inflammation.

Receptors include, but are not limited to, tyrosine kinase receptors which, when activated, result in increased accumulation of PtdIns(3,4,5)P<sub>3</sub>, such as the PDGF receptor, the EGF receptor, members of the FGF receptor family, the CSF-1 receptor, the insulin receptor, the IGF-1 receptor, the SCF (stem cell  
20 factor) receptor, a TNF receptor, such as a TNF- $\alpha$  receptor, and the NGF receptor; receptors associated with the src family non-receptor tyrosine kinases that stimulate PtdIns(3,4,5)P<sub>3</sub> accumulation, such as the IL-2 receptor, IL-3 receptor, mlgM receptor, the CD4 receptor, the CD2 receptor, and the CD3/T cell receptor. Other receptors, such as the cytokine IL-4 receptor and the G  
25 protein linked thrombin receptor, ATP receptor, and the fMLP receptor, that stimulate the activity of a PI3K, resulting in subsequent PtdIns (3,4,5)P<sub>3</sub> accumulation are contemplated herein.

The targeting agents are selected to bind to such cell surface proteins, which must then facilitate internalization of the targeting agent and anything  
30 linked thereto.

The resulting conjugates provided can be used to delivery genes and products to cells that express such receptors. Hence, this provides a means to

-30-

specifically target genes and products to a wide array of cells and in a wide variety of organisms, including plants as well as animals, for effecting genetic therapy and/or delivering products to cells involved in a wide array of disorders and conditions.

5           **2. Preparation of bifunctional molecules**

Methods for preparation of antigen-binding proteins, such as antibodies and antigen-binding fragments thereof that bind to proteins that bind to  $\alpha$  integrins are known to those of skill in the art. The bifunctional molecules can be prepared by recombinant and/or chemical methods. Bifunctional molecules  
10 can be fusion proteins that can be prepared using standard recombinant methods. Bifunctional molecules can be prepared using heterobifunctional reagents and linkers or other suitable chemical conjugation agents. Preparation of bifunctional molecules is exemplified herein, and the exemplified methods can be adapted for preparation of any desired bifunctional molecules.

15           **Plasmids and host cells for expression of constructs encoding bifunctional molecules**

Nucleic acid encoding the selected "P" moiety, such as antibody, generally a heavy chain or portion thereof, is inserted into a suitable vector and expressed in a suitable prokaryotic or eukaryotic host. For antibody expression,  
20 the light chain can be inserted into another plasmid for expression. Numerous suitable hosts and vectors are known and available to those of skill in this art and may be purchased commercially or constructed according to published protocols using well known and available starting materials. Suitable eukaryotic host cells include insect cells, yeast cells, and animal cells. Insect cells and  
25 bacterial host cells are presently preferred. Suitable prokaryotic host cells include E. coli, strains of Bacillus and Streptomyces. For purposes herein, baculovirus expression systems are preferred.

The DNA construct is introduced into a plasmid suitable for expression in the selected host. The sequences of nucleotides in the plasmids that are  
30 regulatory regions, such as promoters and operators, are operationally associated with one another for transcription. The sequence of nucleotides encoding the  $\alpha$  integrin binding protein (designated P) can also include DNA encoding a secretion

-31-

signal, whereby the resulting peptide is a precursor protein. Secretion signals suitable for use are widely available and are well known in the art. Prokaryotic and eukaryotic secretion signals functional in E. coli, may be employed. The presently preferred secretion signals include, but are not limited to, those encoded by the following E. coli genes: ompA, ompT, ompF, ompC, beta-lactamase, pelB and bacterial alkaline phosphatase, and the like (von Heijne (1985) J. Mol. Biol. 184:99-105). In addition, the bacterial pelB gene secretion signal (Lei et al. (1987) J. Bacteriol. 169:4379), the phoA secretion signal, and the cek2 secretion signal, functional in insect cells, may be employed. The most preferred secretion signal for bacterial expression is the E. coli ompA secretion signal. For eukaryotic expression systems, particularly insect cell systems, the signals from secreted proteins, such as insulin, growth hormone, mellitin, and mammalian alkaline phosphatase are of interest herein. Other prokaryotic and eukaryotic secretion signals known to those of skill in the art may also be employed (see, e.g., von Heijne (1985) J. Mol. Biol. 184:99-105). Using the methods described herein, one of skill in the art can substitute secretion signals that are functional in either yeast, insect or mammalian cells to secrete the heterologous protein from those cells. The resulting processed protein may be recovered from the periplasmic space or the fermentation medium or growth medium.

The plasmids can include a selectable marker gene or genes that are functional in the host. A selectable marker gene includes any gene that confers a phenotype on bacteria that allows transformed bacterial cells to be identified and selectively grown from among a vast majority of untransformed cells. Suitable selectable marker genes for bacterial hosts, for example, include the ampicillin resistance gene (Amp<sup>r</sup>), tetracycline resistance gene (Tc<sup>r</sup>) and the kanamycin resistance gene (Kan<sup>r</sup>). The kanamycin resistance gene is presently preferred.

The plasmids used herein preferably include a promoter in operable association with the DNA encoding the protein or polypeptide of interest and are designed for expression of proteins in a bacterial host. It has been found that tightly regulatable promoters are preferred for expression of saporin. Suitable

-32-

promoters for expression of proteins and polypeptides herein are widely available and are well known in the art. For expression of the proteins such promoters are inserted in a plasmid in operative linkage with a control region such as the lac operon. Preferred promoter regions are those that are inducible and

5 functional in E. coli or early genes in vectors of viral origin. Examples of suitable inducible promoters and promoter regions include, but are not limited to: the E. coli lac operator responsive to isopropyl  $\beta$ -D-thiogalactopyranoside (IPTG; see, et al. Nakamura et al. (1979) Cell 18:1109-1117); the metallothionein promoter metal-regulatory-elements responsive to heavy-metal (e.g., zinc) induction (see,

10 e.g., U.S. Patent No. 4,870,009 to Evans et al.); the phage T7lac promoter responsive to IPTG (see, e.g., U.S. Patent No. 4,952,496; and Studier et al. (1990) Meth. Enzymol. 185:60-89) and the TAC promoter. Other promoters include, but are not limited to, the T7 phage promoter and other T7-like phage promoters, such as the T3, T5 and SP6 promoters, the trp, lpp, and lac

15 promoters, such as the lacUV5, from E. coli; the P10 or polyhedrin gene promoter of baculovirus/insect cell expression systems (see, e.g., U.S. Patent Nos. 5,243,041, 5,242,687, 5,266,317, 4,745,051, and 5,169,784) and inducible promoters from other eukaryotic expression systems.

Particularly preferred plasmids for transformation of E. coli cells include

20 the pET expression vectors (see, U.S. patent 4,952,496; available from NOVAGEN, Madison, WI; see, also literature published by Novagen describing the system). Such plasmids include pET 11a, which contains the T7lac promoter, T7 terminator, the inducible E. coli lac operator, and the lac repressor gene; pET 12a-c, which contains the T7 promoter, T7 terminator, and the E. coli

25 ompT secretion signal; and pET 15b (NOVAGEN, Madison, WI), which contains a His-Tag™ leader sequence (Seq. ID NO. 23) for use in purification with a His column and a thrombin cleavage site that permits cleavage following purification over the column; the T7-lac promoter region and the T7 terminator.

Other preferred plasmids include the pKK plasmids, particularly pKK 223-

30 3 (available from Pharmacia; see also, Brosius et al. (1984) Proc. Natl. Acad. Sci. 81:6929; Ausubel et al., Current Protocols in Molecular Biology; U.S. Patent Nos. 5,122,463, 5,173,403, 5,187,153, 5,204,254, 5,212,058,

-33-

5,212,286, 5,215,907, 5,220,013, 5,223,483, and 5,229,279), which contain the TAC promoter. Plasmid pKK has been modified by insertion of a kanamycin resistance cassette with EcoRI sticky ends (purchased from Pharmacia; obtained from pUC4K, see, e.g., Vieira et al. (1982) Gene 19:259-268; and U.S. Patent No. 4,719,179) into the ampicillin resistance marker gene.

Other preferred vectors include the pP<sub>L</sub>-lambda inducible expression vector and the *tac* promoter vector pDR450 (see, e.g., U.S. Patent Nos. 5,281,525, 5,262,309, 5,240,831, 5,231,008, 5,227,469, 5,227,293, ; available from Pharmacia P.L. Biochemicals, see; also Mott, et al. (1985) Proc. Natl. Acad. Sci. U.S.A. 82:88; and De Boer et al. (1983) Proc. Natl. Acad. Sci. U.S.A. 80:21); and baculovirus vectors, such as a pBlueBac vector (also called pJVETL and derivatives thereof; see, e.g., U.S. Patent Nos. 5,278,050, 5,244,805, 5,243,041, 5,242,687, 5,266,317, 4,745,051, and 5,169,784), including pBlueBac III.

Other plasmids include the pIN-IIIompA plasmids (see, U.S. Patent No. 4,575,013 to Inouye; see, also, Duffaud et al. (1987) Meth. Enz. 153:492-507), such as pIN-IIIompA2. The pIN-IIIompA plasmids include an insertion site for heterologous DNA linked in transcriptional reading frame with functional fragments derived from the lipoprotein gene of E. coli. The plasmids also include a DNA fragment coding for the signal peptide of the ompA protein of E. coli, positioned such that the desired polypeptide is expressed with the ompA signal peptide at its amino terminus, thereby allowing efficient secretion across the cytoplasmic membrane. The plasmids further include DNA encoding a specific segment of the E. coli lac promoter-operator, which is positioned in the proper orientation for transcriptional expression of the desired polypeptide, as well as a separate functional E. coli lacI gene encoding the associated repressor molecule that, in the absence of lac operon inducer, interacts with the lac promoter-operator to prevent transcription therefrom. Expression of the desired polypeptide is under the control of the lipoprotein (lpp) promoter and the lac promoter-operator, although transcription from either promoter is normally blocked by the repressor molecule. The repressor is selectively inactivated by

-34-

means of an inducer molecule thereby inducing transcriptional expression of the desired polypeptide from both promoters.

The repressor protein may be encoded by the plasmid containing the construct or a second plasmid that contains a gene encoding for a repressor-  
5 protein. The repressor-protein is capable of repressing the transcription of a promoter that contains sequences of nucleotides to which the repressor-protein binds. The promoter can be derepressed by altering the physiological conditions of the cell. The alteration can be accomplished by the addition to the growth medium of a molecule that inhibits, for example, the ability to interact with the  
10 operator or with regulatory proteins or other regions of the DNA or by altering the temperature of the growth media. Preferred repressor-proteins include, but are not limited to the E. coli lacI repressor responsive to IPTG induction, the temperature sensitive cl857 repressor. The E. coli lacI repressor is preferred.

In certain preferred embodiments, the constructs also include a  
15 transcription terminator sequence. The promoter regions and transcription terminators are each independently selected from the same or different genes. In some embodiments, the DNA fragment is replicated in bacterial cells, preferably in E. coli. The DNA fragment also typically includes a bacterial origin of replication, to ensure the maintenance of the DNA fragment from generation  
20 to generation of the bacteria. In this way, large quantities of the DNA fragment can be produced by replication in bacteria. Preferred bacterial origins of replication include, but are not limited to, the f1-ori and col E1 origins of replication.

Preferred bacterial hosts contain chromosomal copies of DNA encoding  
25 T7 RNA polymerase operably linked to an inducible promoter, such as the lacUV promoter (see, U.S. Patent No. 4,952,496). Such hosts include, but are not limited to, lysogenic E. coli strains HMS174(DE3)pLysS, BL21(DE3)pLysS, HMS174(DE3) and BL21(DE3). Strain BL21(DE3) is preferred. The pLys strains provide low levels of T7 lysozyme, a natural inhibitor of T7 RNA  
30 polymerase. Preferred eukaryotic host are the insect cells Spodoptera frugiperda (sf9 cells; see, e.g., Luckow et al. (1988) Bio/technology 6:47-55 and U.S. Patent No. 4,745,051).

-35-

For insect hosts, which are presently preferred, baculovirus vectors, such as a pTZ (see, e.g., U.S. Patent Nos. 5,278,050, 5,244,805, 5,243,041, 5,242,687, 5,266,317, 4,745,051, and 5,169,784; available from INVITROGEN, San Diego) may also be used for expression of the polypeptides.

- 5 The pBlueBacIII vector is a dual promoter vector and provides for the selection of recombinants by blue/white screening as this plasmid contains the  $\beta$ -galactosidase gene (lacZ) under the control of the insect recognizable ETL promoter and is inducible with IPTG. A DNA construct is introduced into a baculovirus vector pBluebac III (INVITROGEN, San Diego, CA) and then co-
- 10 transfected with wildtype virus into insect cells Spodoptera frugiperda (sf9 cells; see, e.g., Luckow et al. (1988) Bio/technology 6:47-55 and U.S. Patent No. 4,745,051).

- Other baculovirus vectors, such as pPbac and pMbac (available from Stratagene, San Diego, CA, see, also Lernhardt et al. (1993) Strategies 6:20-21,
- 15 and the Stratagene Catalog page 218), which contain the human alkaline phosphatase (see, e.g., Bailey et al. (1989) Proc. Natl. Acad. Sci. U. S. A. 86:22-26) and melittin (see, e.g., Tessier et al. (1991) Gene 98:177-183) secretory signals inserted into the BamHI and NdeI sites, respectively of pJVP10Z (see, e.g., Kawamoto et al. (1991) Biochem. Biophys. Res. Commun.
- 20 181:756-63, Ueda et al. (1994) Gene 140:267-272, are also suitable for use herein, particularly if secretion is desired. Insertion of genes into the SmaI/BamHI sites of these vectors results in fusion proteins that are directed into the insect cell secretory pathway, which processes the pro-polypeptide so that mature peptide or fusion protein is secreted into the growth medium. Other
- 25 heterologous signal sequences, such as the insulin signal sequence (see, e.g., U.S. Patent No. 4,431,746 for DNA encoding the signal sequence), the growth hormone signal sequence, mammalian alkaline phosphatase, the mellitin signal sequence and others that are processed by insect cells are used.

- The constructs provided herein have can be inserted into the baculovirus
- 30 vector sold commercially under the name pBLUEBACIII (INVITROGEN, San Diego CA; see the INVITROGEN CATALOG; see, Vialard et al. (1990) J. Virol. 64:37; see also, U.S. Patent No. 5,270,458; U.S. Patent No. 5,243,041; and



-36-

published International PCT Application WO 93/10139, which is based on U.S. patent application Serial No. 07/792,600. The pBlueBacIII vector is a dual promoter vector and provides for the selection of recombinants by blue/white screening as this plasmid contains the  $\beta$ -galactosidase gene (lacZ) under the control of the insect recognizable ETL promoter and is inducible with IPTG. The construct of interest is inserted into this vector under control of the polyhedrin promoter. The DNA is then cotransfected, such as by  $\text{Ca}(\text{PO}_4)_2$  or calcium phosphate transfection or liposomes, into Spodoptera frugiperda cells (sf9 cells) with wildtype baculovirus and grown in tissue culture flasks or in suspension cultures. Blue occlusion minus viral plaques are selected and plaque purified and screened for expression. Details are set forth in the Examples.

#### Combinatorial methods

Molecules of desired specificity can be prepared using a variety of protocols in which portions of antibody molecules are combined to produce antigen binding molecules and are screened for desired specificity. Generally for phage display libraries, at least the Fab fragment is required, since secretion of the heterodimeric antibody fragment is involved.

Complementarily determining regions (CDRs) that are present in the Fab fragment can be varied by random mutagenesis (creation of "synthetic antibody libraries") or by other methods. A phagemid vector such as pCombIII, designed to express the heavy chain antibody fragment as an N-terminal fusion with the pIII phage coat protein domain (Barbas *et al.* (1991) *Proc. Natl. Acad. Sci. USA*, 88:7978) can be used for heterodimeric expression of the library of Fab antibody fragments (or Fab'2 antibody fragments) on the surface of phage, using bacteria.

25

Thus, for example, the DAV-1 heavy chain fused to the growth factor or a portion of the growth factor involved in binding to its receptor may be used to form the phage display library, and the library may then be "panned" for bifunctional antibodies that bind with high specificity to both the adenoviral surface protein and to the growth factor/cytokine receptor.

30

-37-

### Linkers and chemical conjugation

For fusion protein the linkers are peptides; for chemically conjugated molecules the linkers may be other moieties. In addition, a combinations of chemically conjugated molecules and recombinantly produced portions, such as

5 recombinantly produced antibodies or fragments or other antigen binding molecules may be combined and chemically fused to targeting agents, such as a growth factor or hormone. If a linker is used it is selected such that it does not interfere with the activity of the targeted agent upon interaction of the conjugate with a cell surface protein. Any appropriate linker known to those of skill in this

10 art may be used. The linker may be selected to improve activity by permitting the targeted agent to complex with the viral or bacterial vector. In some instances the linker is selected to increase the specificity, toxicity, solubility, serum stability, and/or intracellular availability targeted moiety. In some

15 embodiments, several linkers may be included in order to take advantage of desired properties of each linker. Flexible linkers and linkers that increase solubility of the conjugates are contemplated for use, either alone or with other linkers are contemplated herein.

For chemical conjugation the components of the bifunctional molecules may be directly linked or attached via a linker. Linkers that are suitable for

20 chemically linking conjugates include, but are not limited to, disulfide bonds, thioether bonds, hindered disulfide bonds, esters, and covalent bonds between free reactive groups, such as amine and thiol groups. These bonds are produced using heterobifunctional reagents to produce reactive thiol groups on one or both

25 of the polypeptides and then reacting the thiol groups on one polypeptide with reactive thiol groups or amine groups on the other. Other linkers include, but are not limited to: acid cleavable linkers, such as bismaleimidoethoxy propane; acid labile-transferrin conjugates and adipic acid dihydrazide that are cleaved in more

acidic environments; photocleavable cross linkers that are cleaved by visible or UV light.

30 Linkers that are suitable for chemically linked conjugates include, but are not limited to: disulfide bonds; thioether bonds; hindered disulfide bonds; and covalent bonds between free reactive groups, such as amine and thiol groups.

-38-

These bonds are produced using heterobifunctional reagents to produce reactive thiol groups on one or both of the polypeptides and then reacting the thiol groups on one polypeptide with reactive thiol groups or amine groups to which reactive maleimido groups or thiol groups can be attached on the other. Other

5 linkers include, acid cleavable linkers, such as bismaleimidoethoxy propane, acid labile-transferrin conjugates and adipic acid dihydrazide, that would be cleaved in more acidic intracellular compartments; cross linkers that are cleaved upon exposure to UV or visible light; and linkers, such as the various domains, such as C<sub>H</sub>1, C<sub>H</sub>2, and C<sub>H</sub>3, from the constant region of human IgG, (see, Batra *et al.*

10 (1993) *Molecular Immunol.* 30:379-386). In some embodiments, several linkers may be included in order to take advantage of desired properties of each linker.

Chemical linkers and peptide linkers may be inserted by covalently coupling the linker to the TA and the P portion (the  $\alpha$  integrin protein binding

15 portion, such as an antibody). The heterobifunctional agents, described below, may be used to effect such covalent coupling. Peptide linkers may also be linked by expressing DNA encoding the linker and TA, linker and P, or linker, targeted agent and TA as a fusion protein.

Numerous heterobifunctional cross-linking reagents that are used to form

20 covalent bonds between amino groups and thiol groups and to introduce thiol groups into proteins, are known to those of skill in this art (see, *e.g.*, the PIERCE CATALOG, ImmunoTechnology Catalog & Handbook, 1992-1993, which describes the preparation of and use of such reagents and provides a commercial source for such reagents; see, also, *e.g.*, Cumber *et al.* (1992) *Bioconjugate*

25 *Chem.* 3:397-401; Thorpe *et al.* (1987) *Cancer Res.* 47:5924-5931; Gordon *et al.* (1987) *Proc. Natl. Acad. Sci.* 84:308-312; Walden *et al.* (1986) *J. Mol. Cell Immunol.* 2:191-197; Carlsson *et al.* (1978) *Biochem. J.* 173: 723-737; Mahan *et al.* (1987) *Anal. Biochem.* 162:163-170; Wawryznaczk *et al.* (1992) *Br. J. Cancer* 66:361-366; Fattom *et al.* (1992) *Infection & Immun.* 60:584-589).

30 These reagents may be used to form covalent bonds between the TA and targeted agent. These reagents include, but are not limited to: N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP; disulfide linker); sulfosuccinimidyl 6-[3-(2-

-39-

pyridyldithio)propionamido]hexanoate (sulfo-LC-SPDP); succinimidyloxycarbonyl- $\alpha$ -methyl benzyl thiosulfate (SMBT, hindered disulfate linker); succinimidyl 6-[3-(2-pyridyldithio) propionamido]hexanoate (LC-SPDP); sulfosuccinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate (sulfo-SMCC); succinimidyl 3-(2-pyridyldithio)butyrate (SPDB; hindered disulfide bond linker); sulfosuccinimidyl 2-(7-azido-4-methylcoumarin-3-acetamide) ethyl-1,3'-dithiopropionate (SAED); sulfo-succinimidyl 7-azido-4-methylcoumarin-3-acetate (SAMCA); sulfosuccinimidyl 6-[ $\alpha$ -methyl- $\alpha$ -(2-pyridyldithio)toluamido]hexanoate (sulfo-LC-SMPT); 1,4-di-[3'-(2'-pyridyldithio)propionamido]butane (DPDPB); 4-succinimidyloxycarbonyl- $\alpha$ -methyl- $\alpha$ -(2-pyridyldithio)toluene (SMPT, hindered disulfate linker); sulfo-succinimidyl 6-[ $\alpha$ -methyl- $\alpha$ -(2-pyridyldithio)toluamido]hexanoate (sulfo-LC-SMPT); *m*-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS); *m*-maleimidobenzoyl-N-hydroxysulfosuccinimide ester (sulfo-MBS); N-succinimidyl(4-iodoacetyl)aminobenzoate (SIAB; thioether linker); sulfosuccinimidyl(4-iodoacetyl)amino benzoate (sulfo-SIAB); succinimidyl 4-(*p*-maleimidophenyl)butyrate (SMPB); sulfosuccinimidyl 4-(*p*-maleimidophenyl)butyrate (sulfo-SMPB); azidobenzoyl hydrazide (ABH).

Acid cleavable linkers, photocleavable and heat sensitive linkers may also be used, particularly where it may be necessary to cleave the targeted agent to permit it to be more readily accessible to reaction.

Acid cleavable linkers include, but are not limited to, bismaleimidoethoxy propane; and adipic acid dihydrazide linkers (see, e.g., Fattom et al. (1992) Infection & Immun. 60:584-589) and acid labile transferrin conjugates that contain a sufficient portion of transferrin to permit entry into the intracellular transferrin cycling pathway (see, e.g., Welhöner et al. (1991) J. Biol. Chem. 266:4309-4314).

Photocleavable linkers are linkers that are cleaved upon exposure to light (see, e.g., Goldmacher et al. (1992) Bioconj. Chem. 3:104-107, which linkers are herein incorporated by reference), thereby releasing the targeting agent from the bifunctional molecule and the complex upon exposure to light. Photocleavable linkers that are cleaved upon exposure to light are known (see, e.g., Hazum et al. (1981) in Pept., Proc. Eur. Pept. Symp., 16th, Brunfeldt, K

-40-

- (Ed), pp. 105-110, which describes the use of a nitrobenzyl group as a photocleavable protective group for cysteine; Yen et al. (1989) Makromol. Chem 190:69-82, which describes water soluble photocleavable copolymers, including hydroxypropylmethacrylamide copolymer, glycine copolymer, fluorescein
- 5 copolymer and methylrhodamine copolymer; Goldmacher et al. (1992) Bioconj. Chem. 3:104-107, which describes a cross-linker and reagent that undergoes photolytic degradation upon exposure to near UV light (350 nm); and Senter et al. (1985) Photochem. Photobiol 42:231-237, which describes
- 10 nitrobenzyloxycarbonyl chloride cross linking reagents that produce photocleavable linkages), thereby releasing the targeted agent upon exposure to light. Such linkers would have particular use in treating dermatological or ophthalmic conditions that can be exposed to light using fiber optics. After administration of the conjugate, the eye or skin or other body part can be exposed to light, resulting in release of the targeted moiety from the conjugate.
- 15 Such photocleavable linkers are useful in connection with diagnostic protocols in which it is desirable to remove the targeting agent to permit rapid clearance from the body of the animal.

3. Preparation of complexes of the bifunctional molecules with viral vector particles or bacterial particles

- 20 The complexes between the bifunctional molecules and vectors can be prepared by incubating the vectors and bifunctional molecules under suitable conditions to effect formation of the complexes.

Assays for activity

- Any assay to assess the ability of the resulting complexes to
- 25 deliver genes. Such assays are known to those of skill in the art, and several are exemplified herein (see EXAMPLES).

4. Exemplary embodiment DAV-1 constructs and use thereof for targeting

- Exemplary of the embodiments contemplated herein are
- 30 bifunctional molecules formed by linkage of a targeting agent, such as a growth factor known to bind to receptors that activate PI3K, to an antibody or antigen-binding portion thereof, that specifically binds to the penton protein of a variety of adenovirus serotypes. It is understood that these embodiments are exemplary

-41-

and that any antibody that binds to a protein on a viral particle or bacterial cell or other moiety intended for delivery into a cell can be used to specifically target such moieties to cells by virtue of interaction of the targeting agent with cell surface receptors. Presently preferred are adenovirus particles and the penton protein thereof that is responsible for interaction with  $\alpha v$  integrins, which promote viral internalization.

A penton base monoclonal antibody, DAV-1, which recognizes the integrin binding site on adenovirus particle types 2/5 (Stewart *et al.* (1997) *EMBO J.* 16:1189-1198; EXAMPLE 1), binds to the penton base with high affinity but does not inhibit virus infection is re-engineered herein (see, EXAMPLE 1). Nucleic acid encoding the DAV-1 heavy chain (see, *e.g.*, SEQ ID Nos. 5) has been fused with one of several different cytokines/growth factors known to activate PI3K and the encoded fusion proteins are co-expressed in insect cells with the DAV-1 light chain. The resulting bifunctional molecules retain immunoreactivity and cytokine function.

The bifunctional molecules have been complexed with adenovirus and shown to provide increased gene delivery to cells lacking  $\alpha v$  integrins, as well as cells expressing  $\alpha v$  integrins, evidencing PI3K activation. Complexing with the bifunctional molecule also allows gene delivery by a fiberless Ad vector that lacks the ability to bind CAR (see, EXAMPLES 5 and 6).

a. Analysis of bifunctional signaling antibodies

DAV-1 bifunctional signaling antibodies, designated DT (DAV-1 fused to TNF- $\alpha$ ), DI (DAV-1 fused to IGF-1), and DE (DAV-1 fused to EGF), were expressed in insect cells as secreted proteins and purified on Protein L affinity columns. The DT heavy chain ( $D_H T$ ) had an apparent molecular weight of approximately 70 kDa, consistent with the combined sizes of the DAV-1  $\gamma$  heavy chain (53 kDa) and monomeric TNF- $\alpha$  ligand (17 kDa). The apparent molecular weight of the  $\kappa$  light chains of DAV-1 ( $D_L$ ) was identical to that of the recombinant DT molecule (approx. 25 kDa). Western blot analyses showed that the DAV-1 mAb (D) and the DT molecules were recognized by an anti-mouse IgG antibody, while only the DT molecule was recognized by an anti-TNF- $\alpha$  polyclonal antibody.

-42-

DT molecules were capable of binding to immobilized penton base or Ad particles in an ELISA and elicited cytotoxicity against a TNF- $\alpha$  sensitive cell line, MCF-7, indicating that the DT bifunctional molecule retains virus and cytokine receptor binding functions.

5                                    b.        Bifunctional molecules promote Ad-mediated gene delivery to cells lacking  $\alpha v$  integrins

A first generation adenovirus vector containing a RSV-driven *LacZ* reporter gene with DT was preincubated at a ratio of 2 antibody molecules per RGD motif. This complex was then added to M21-L12 human melanoma cells, which do not express  $\alpha v$  integrins (Felding-Habermann *et al.* (1992) *J.Clin.Invest.* 89:2018-2022), but can support efficient virus binding (Wickham *et al.* (1993) *Cell* 73:309-319). Ad complexed with DT but not D alone, significantly increases Ad-mediated gene delivery to M21-L12 cells as measured by transgene expression at 48 hrs post-infection. Approximately 60% of cells incubated with Ad plus DT stained positive for  $\beta$ -galactosidase, compared to less than 3% of cells that had been incubated with virus alone or virus plus D. The increase in gene delivery by DT was not due to increased activation of the RSV LTR transgene promoter as a consequence of ligation of the TNF receptor, since M21-L12 cells that had been infected with adenovirus alone for three hours followed by addition of DT showed very little increase in gene delivery at 48 hours postinfection. This result indicates that bifunctional molecules increase adenovirus-mediated gene delivery by enhancing one or more steps associated with cell entry.

                                  c.        DT molecules enhance Ad binding and internalization

25        The following experiments showed that DT enhancement of gene delivery was associated with increased virus attachment to M21-L12 cells. Pre-incubation of  $^{125}$ I-labeled Ad particles with DT but not D molecules increased virus binding approximately 5 fold. To investigate the molecules responsible for increased binding, competition experiments were performed. Ad-DT binding to cells was measured in the presence of a 50-fold excess of recombinant fiber protein or anti-TNF- $\alpha$  or a combination of these molecules. Either recombinant fiber or anti-TNF- $\alpha$  antibody alone was capable of blocking only 20-25% of Ad-

30

-43-

DT binding to cells. In contrast, approximately 70% of binding could be inhibited by a combination of fiber and anti-TNF- $\alpha$ . These findings indicate that Ad-DT binding to cells is mediated by CAR-fiber interaction as well as TNF- $\alpha$  receptor association.

- 5                   d.     DT molecules potentiate internalization of  $^{125}$ I-labeled virus particles as measured by resistance to trypsin digestion.

As demonstrated in earlier studies(Wickham *et al.* (1993) *Cell* 73:309-319), M21-L12 cells support relatively low levels of adenovirus internalization. This is due to the absence of  $\alpha$ v integrins. DT molecules significantly increased  
10 the rate and extent of adenovirus internalization into these cells. Together, these findings indicate that DT molecules enhance gene delivery by promoting virus binding as well as virus internalization.

- e.     DT enhancement of gene delivery is associated with PI3K activation

- 15               Efficient Ad internalization via  $\alpha$ v integrins requires activation of PI3K, a key cellular signaling molecule. M21-L12 cells were treated with PI3K inhibitors, wortmannin or LY292004 prior to virus infection to show that DT enhancement of gene delivery also involves PI3K. Wortmannin and LY294002 inhibited Ad-mediated gene delivery by approximately 70% and 50%,  
20 respectively, indicating that PI3K activity plays a major role in DT enhancement of gene delivery.

- For further demonstration of the role of PI3K-dependent signaling in enhanced gene delivery, Ad-mediated gene delivery by other bifunctional molecules whose cytokine/growth factor domains are known to activate PI3K  
25 was measured. DT, DE and DI molecules enhanced gene delivery by approximately 30, 10 and 5 fold respectively. Enhanced gene delivery by these molecules was also inhibited by pretreatment of cells with wortmannin. These findings further demonstrate that PI3K activation promotes Ad gene delivery.



-44-

f. Bifunctional molecules allow gene delivery by fiberless adenovirus particles

Fiberless adenovirus vectors that cannot bind to CAR have been constructed (see, *e.g.*, copending U.S. application Serial No. 09/482,682 and  
5 U.S. application Serial No. 09/562,934; see, also Von Seggern *et al.* (1999) *J. Virol.* 73:1601-1608). The structure of these particles is nearly identical to that of wildtype virions. Fiberless particles alone showed almost no transgene delivery to SW480 epithelial cells, even though these cells express both CAR and integrin  $\alpha\beta 5$  (Von Seggern *et al.* (1999) *J. Virol.* 73:1601-1608).

10 DT molecules enhanced gene delivery of fiberless viruses. Fiberless particles complexed with DT or DI molecules exhibited increased gene delivery approximately 10-15, 3 and 5 fold, respectively, compared to the uncomplexed fiberless particles. These findings indicate that fiberless particles can be retargeted to cells via signal transducing antibodies.

15 The use of fiberless adenoviral vectors for gene therapy is desirable because it is then possible to restrict viral tropism to selected cell types by adopting a specific targeting strategy and abrogating the interaction between the viral fiber protein and the CAR receptors.

g. Summary of results

20 It is demonstrated herein that Ad vectors complexed with bifunctional molecules significantly increased gene delivery to human melanoma cells lacking  $\alpha\upsilon$  integrins and was associated with both increased virus binding and internalization compared to Ad vectors without such bifunctional molecules. Importantly, PI3K activation plays a key role in this process as a significant  
25 reduction in delivery following treatment of cells with pharmacologic inhibitors of PI3K was observed. Moreover, bifunctional molecules displaying distinct growth factor ligands, each of which is known to activate PI3K, also enhanced gene delivery. The variability in gene delivery by different bifunctional molecules is most likely to be due to differences in the expression of growth factor/cytokine  
30 receptors on different cell types or to the extent of PI3K activation induced by each ligand.

-45-

Achieving gene delivery to specific cell types has been hampered by the fiber receptor (CAR), which is expressed on multiple cell types. To circumvent this problem, a fiberless adenovirus vector was retargeted with bifunctional molecules. Significant enhancement of gene delivery by a fiberless adenovirus  
5 complexed with several different bifunctional molecules was detected. These studies confirm the structural and functional integrity of fiberless adenovirus particles and indicate their usefulness for retargeting in gene delivery.

The PI3K-dependent signaling pathway should be useful for increasing the uptake of other viral vectors. For example, adeno-associated virus (AAV) has  
10 also been reported to use  $\alpha v$  integrins for infection. Because a variety of bacterial and viral pathogens use integrins and/or PI3K activation for host cell invasion indicates that targeting of PI3K-dependent signaling pathways can also be used as a general scheme for potentiating cell entry of nonviral vectors. The exemplified methods and bifunctional molecules can be adapted for use for  
15 potentiating entry of other viral vectors and bacterial vectors that use the PI3K-dependent signaling pathways for internalization.

C. Construction of the viral particles

1. Selection of viral genome and fiber protein

Methods for preparing recombinant adenoviral vectors for gene  
20 product delivery are well known. Preferred among those are the methods exemplified herein (see EXAMPLES) and also described in copending U.S. application Serial No. 09/482,682 (also filed as International PCT application No. PCT/US00/00265, filed January 14, 2000, which claims priority to U.S. provisional application Serial No. 60/115,920, as does U.S. application Serial  
25 No. 09/482,682)).

As noted, any desired recombinant adenovirus is contemplated for use in the methods herein as long as the viral genome is packaged in a capsid that includes at least the portion of a fiber protein that provides selective binding to photoreceptor cells. This fiber protein is preferably from an adenovirus type D  
30 serotype and is preferably an Ad37 fiber. The fiber protein should retain the knob region at the C-terminus ("head domain") from the Ad type D virus that contains the type-specific antigen and is responsible for binding to the cell

-46-

surface receptor. Hence the fiber protein can be a chimeric fiber protein as long as it retains a sufficient portion of the type-D serotype to specifically bind to photoreceptor cells. Generally the portion retained will be all or a portion of the knob region. The precise amount of knob region required can be determined empirically by including portions thereof and identifying the minimum residues from and Ad type D serotype, preferably Ad37, to effect selective targeting of a virion packaged with such fiber to photoreceptors in the eye upon introduction of the packaged virion into the aqueous humor.

Recombinant adenovirus containing heterologous nucleic acids that encode a desired product, such a gene to correct a genetic defect, may be made by any methods known to those of skill in the art. The viruses are packaged in a suitable cell line.

The family of Adenoviridae includes many members with at least 51 known serotypes of human adenovirus (Ad1-Ad47) (Shenk, *Virology*, Chapter 67, in Fields *et al.*, eds. Lippincott-Raven, Philadelphia, 1996,) as well as members of the genus Mastadenovirus including human, simian, bovine, equine, porcine, ovine, canine and opossum viruses, and members of the Aviadenovirus genus, including bird viruses, e.g. CELO. Thus it is contemplated that the methods herein can be applied to any recombinant viral vectors derived from any adenovirus species. One of skill in the art would have knowledge of the different adenoviruses (see, e.g., Shenk, *Virology*, Chapter 67, in Fields *et al.*, eds. Lippincott-Raven, Philadelphia, 1996,) and can construct recombinant viruses containing portions of the genome of any such virus. The methods herein may also be adapted for use with any delivery vehicle that internalizes via receptors that use the PI3K signalling pathway, particularly that employ  $\alpha$  integrins in the process.

## 2. Packaging

Recombinant adenoviral vectors generally have at least a deletion in the first viral early gene region, referred to as E1, which includes the E1a and E1b regions. Deletion of the viral E1 region renders the recombinant adenovirus defective for replication and incapable of producing infectious viral particles in subsequently-infected target cells. Thus, to generate E1-deleted adenovirus

-47-

genome replication and to produce virus particles requires a system of complementation which provides the missing E1 gene product. E1 complementation is typically provided by a cell line expressing E1, such as the human embryonic kidney packaging cell line, i.e. an epithelial cell line; called

5 293. Cell line 293 contains the E1 region of adenovirus, which provides E1 gene region products to "support" the growth of E1-deleted virus in the cell line (see, e.g., Graham *et al.*, *J. Gen. Virol.* 36: 59-71, 1977). Additionally, cell lines that may be usable for production of defective adenovirus having a portion of the adenovirus E4 region have been reported (WO 96/22378). Multiply deficient

10 adenoviral vectors and complementing cell lines have also been described (WO 95/34671, U.S. Patent No. 5,994,106).

Copending U.S. application Serial No. 09/482,682 (also filed as International PCT application No. PCT/US00/00265, filed January 14, 2000)) provides packaging cell lines that support viral vectors with deletions of major

15 portions of the viral genome, without the need for helper viruses and also provides cell lines and helper viruses for use with helper-dependent vectors.

The copending application provides a packaging cell line that has heterologous DNA stably integrated into the chromosomes of the cellular genome. The heterologous DNA sequence encodes one or more adenovirus

20 regulatory and/or structural polypeptides that complement the genes deleted or mutated in the adenovirus vector genome to be replicated and packaged. The packaging cell line expresses, for example, one or more adenovirus structural proteins, polypeptides, or fragments thereof, such as penton base, hexon, fiber, polypeptide IIIa, polypeptide V, polypeptide VI, polypeptide VII, polypeptide VIII,

25 and biologically active fragments thereof. The expression can be constitutive or under the control of a regulatable promoter. These cell lines are particularly designed for expression of recombinant adenoviruses intended for delivery of therapeutic products.

Particular packaging cell lines complement viral vectors having a deletion

30 or mutation of a DNA sequence encoding an adenovirus structural protein, regulatory polypeptides E1A and E1B, and/or one or more of the following regulatory proteins or polypeptides: E2A, E2B, E3, E4, L4, or fragments thereof.

-48-

The packaging cell lines are produced by introducing each DNA molecule into the cells and then into the genome via a separate complementing plasmid or plurality of DNA molecules encoding the complementing proteins can be introduced via a single complementing plasmid.

- 5 For therapeutic applications, the delivery plasmid further includes a nucleotide sequence encoding a foreign polypeptide. Exemplary delivery plasmids is pDV44, pE1B gal and pE1sp1B. In a similar or analogous manner, therapeutic genes may be introduced.

- 10 The cell further includes a complementing plasmid encoding a fiber as contemplated herein; the plasmid or portion thereof is integrated into a chromosome(s) of the cellular genome of the cell.

- 15 In one embodiment, a composition comprises a cell containing first and second delivery plasmids wherein a first delivery plasmid comprises an adenovirus genome lacking a nucleotide sequence encoding fiber and incapable of directing the packaging of new viral particles in the absence of a second delivery plasmid, and a second delivery plasmid comprises an adenoviral genome capable of directing the packaging of new viral particles in the presence of the first delivery plasmid.

- 20 The packaging cell line can be derived from a procaryotic cell line or from a eukaryotic cell line. While various embodiments suggest the use of mammalian cells, and more particularly, epithelial cell lines, a variety of other, non-epithelial cell lines are used in various embodiments.

### 3. Components of the nucleic acid molecule included in the particle

- 25 A recombinant viral vector or therapeutic viral vector for use in the methods herein, typically includes a nucleic acid fragment that encodes a protein or polypeptide molecule, or a biologically active fragment thereof, or other regulatory sequence, that is intended for use for therapeutic applications.

- 30 The nucleic acid molecule to be packaged in the viral particle also may include an enhancer element and/or a promoter located 3' or 5' to and controlling the expression of the therapeutic product-encoding nucleic acid molecule if the product is a protein. Further, for purposes herein, the promoter and/or other transcriptional and translational regulatory sequences controlling

-49-

expression of the product is preferably one that is expressed specifically in the targeted cells, such as the a photoreceptor-specific promoter, such as a rhodopsin gene promoter.

The nucleic acid molecule to be packaged in viral capsid includes at least  
5 2 different operatively linked DNA segments. The DNA can be manipulated and amplified by PCR as described herein and by using standard techniques, such as those described in *Molecular Cloning: A Laboratory Manual, 2nd Ed.*, Sambrook *et al.*, eds., Cold Spring Harbor, New York (1989). Typically, to produce such molecule, the sequence encoding the selected polypeptide and the promoter or  
10 enhancer are operatively linked to a DNA molecule capable of autonomous replication in a cell either *in vivo* or *in vitro*. By operatively linking the enhancer element or promoter and nucleic acid molecule to the vector, the attached segments are replicated along with the vector sequences.

Thus, the recombinant DNA molecule (rDNA) is a hybrid DNA molecule  
15 comprising at least 2 nucleotide sequences not normally found together in nature. In various preferred embodiments, one of the sequences is a sequence encoding an Ad-derived polypeptide, protein, or fragment thereof. The nucleic acid molecule intended to be packaged is from about 20 base pairs to about 40,000 base pairs in length, preferably about 50 bp to about 38,000 bp in  
20 length. In various embodiments, the nucleic acid molecule is of sufficient length to encode one or more adenovirus proteins or functional polypeptide portions thereof. Since individual Ad polypeptides vary in length from about 19 amino acid residues to about 967 amino acid residues, encoding nucleic acid molecules from about 50 bp up to about 3000 bp, depending on the number and size of  
25 individual polypeptide-encoding sequences that are "replaced" in the viral vectors by therapeutic product-encoding nucleic acid molecules.

Preferably the molecule includes an adenovirus tripartite leader (TPL) nucleic acid sequence operatively linked to an intron containing RNA processing signals (such as for example, splice donor or splice acceptor sites) suitable for  
30 expression in the packaging cell line. Most preferably the intron contains a splice donor site and a splice acceptor site. Alternatively, the TPL nucleotide sequence may not comprise an intron. The intron includes any sequence of

-50-

nucleotides that function in the packaging cell line to provide RNA processing signals, including splicing signals. Introns have been well characterized from a large number of structural genes, and include but are not limited to a native intron 1 from adenovirus, such as Ad5's TPL intron 1; others include the SV40  
5 VP intron; the rabbit beta-globin intron, and synthetic intron constructs (see, e.g., Petitclerc *et al.* (1995) *J. Biothechnol.*, 40:169; and Choi *et al.* (1991 *Mol. Cell. Biol.*, 11:3070).

The nucleic acid molecule encoding the TPL includes either (a) first and second TPL exons or (b) first, second and third TPL exons, where each TPL exon  
10 in the sequence is selected from among the complete TPL exon 1, partial TPL exon 1, complete TPL exon 2 and complete TPL exon 3. A complete exon is one which contains the complete nucleic acid sequence based on the sequence found in the wildtype viral genome. Preferably the TPL exons are from Ad2, Ad3, Ad5, Ad7 and the like, however, they may come from any Ad serotype, as  
15 described herein. A preferred partial TPL exon 1 is described in the Examples. The use of a TPL with a partial exon 1 has been reported (International PCT application No. WO 98/13499).

The intron and the TPL exons can be operatively linked in a variety of configurations to provide a functional TPL nucleotide sequence. An intron may  
20 not be a part of the construct. For example, the intron can be positioned between any of TPL exons 1, 2 or 3, and the exons can be in any order of first and second, or first/second/third. The intron can also be placed preceding the first TPL exon or following the last TPL exon. In a preferred embodiment, complete TPL exon 1 is operatively linked to complete TPL exon 2 operatively  
25 linked to complete TPL exon 3. In a preferred variation, adenovirus TPL intron 1 is positioned between complete TPL exon 1 and complete TPL exon 2. It may also be possible to use analogous translational regulators from other viral systems such as rabiesvirus.

#### 4. Complementing Plasmids

30 Also contemplated are the use of nucleic acid molecules, typically in the form of DNA plasmid vectors, which are capable of expression of an adenovirus structural protein or regulatory protein. Because these expression plasmids are

-51-

used to complement the defective genes of a recombinant adenovirus vector genome, the plasmids are referred to as complementing or complementation plasmids.

The complementing plasmid contains an expression cassette, a nucleotide  
5 sequence capable of expressing a protein product encoded by the nucleic acid molecule. Expression cassettes typically contain a promoter and a structural gene operatively linked to the promoter. The complementing plasmid can further include a sequence of nucleotides encoding TPL nucleotide to enhance  
10 expression of the structural gene product when used in the context of adenovirus genome replication and packaging.

A complementing plasmid can include a promoter operatively linked to a sequence of nucleotides encoding an adenovirus structural polypeptide, such as, but are not limited to, penton base; hexon; fiber; polypeptide IIIa; polypeptide V; polypeptide VI; polypeptide VII; polypeptide VIII; and biologically active  
15 fragments thereof. In another variation, a complementing plasmid may also include a sequence of nucleotides encoding a first adenovirus regulatory polypeptide, a second regulatory polypeptide, and/or a third regulatory polypeptide; or any combination of the foregoing.

#### 5. Nucleic Acid Molecule Synthesis

20 A nucleic acid molecule comprising synthetic oligonucleotides can be prepared using any suitable method, such as, the phosphotriester or phosphodiester methods (see, e.g., Narang (1979) *et al.*, *Meth. Enzymol.*, 68:90; U.S. Patent No. 4,356,270; and Brown *et al.*, (1979) *Meth. Enzymol.*, 68:109). For oligonucleotides, the synthesis of the family members can be  
25 conducted simultaneously in a single reaction vessel, or can be synthesized independently and later admixed in preselected molar ratios. For simultaneous synthesis, the nucleotide residues that are conserved at preselected positions of the sequence of the family member can be introduced in a chemical synthesis protocol simultaneously to the variants by the addition of a single preselected  
30 nucleotide precursor to the solid phase oligonucleotide reaction admixture when that position number of the oligonucleotide is being chemically added to the growing oligonucleotide polymer. The addition of nucleotide residues to those



-52-

- positions in the sequence that vary can be introduced simultaneously by the addition of amounts, preferably equimolar amounts, of multiple preselected nucleotide precursors to the solid phase oligonucleotide reaction admixture during chemical synthesis. For example, where all four possible natural
- 5 nucleotides (A,T,G and C) are to be added at a preselected position, their precursors are added to the oligonucleotide synthesis reaction at that step to simultaneously form four variants (see, e.g., Ausubel *et al.* (*Current Protocols in Molecular Biology*, Suppl. 8. p.2.11.7, John Wiley & Sons, Inc., New York ,1991).
- 10 Nucleotide bases other than the common four nucleotides (A,T,G or C), or the RNA equivalent nucleotide uracil (U), can also be used. For example, it is well known that inosine (I) is capable of hybridizing with A, T and G, but not C. Examples of other useful nucleotide analogs are known in the art and may be found referred to in 37 C.F.R. §1.822.
- 15 Thus, where all four common nucleotides are to occupy a single position of a family of oligonucleotides, that is, where the preselected nucleotide sequence is designed to contain oligonucleotides that can hybridize to four sequences that vary at one position, several different oligonucleotide structures are contemplated. The composition can contain four members, where a
- 20 preselected position contains A,T,G or C. Alternatively, a composition can contain two nucleotide sequence members, where a preselected position contains I or C, and has the capacity the hybridize at that position to all four possible common nucleotides. Finally, other nucleotides may be included at the preselected position that have the capacity to hybridize in a non-de stabilizing
- 25 manner with more than one of the common nucleotides in a manner similar to inosine.

Similarly, larger nucleic acid molecules can be constructed in synthetic oligonucleotide pieces, and assembled by complementary hybridization and ligation, as is well known.

30 D. Adenovirus Expression Vector Systems

The adenovirus vector genome that is encapsulated in the virus particle and that expresses exogenous genes in a gene therapy setting is a key

-53-

component of the system, which systems are well known and readily available. Thus, the components of a recombinant adenovirus vector genome include the ability to express selected adenovirus structural genes, to express a desired exogenous protein, and to contain sufficient replication and packaging signals  
5 that the genome is packaged into a gene delivery vector particle. The preferred replication signal is an adenovirus inverted terminal repeat containing an adenovirus origin of replication, as is well known.

Although adenovirus include many proteins, not all adenovirus proteins are required for assembly of a recombinant adenovirus particle (vector). Thus,  
10 deletion of the appropriate genes from a recombinant Ad vector permits accommodation of even larger "foreign" DNA segments.

Particularly contemplated are helper dependent systems as described above, in which the adenovirus vector genome does not encode a functional adenovirus fiber protein. A non-functional fiber gene refers to a deletion,  
15 mutation or other modification to the adenovirus fiber gene such that the gene does not express any or insufficient adenovirus fiber protein to package a fiber-containing adenovirus particle without complementation of the fiber gene by a complementing plasmid or packaging cell line. Such a genome is referred to as a "fiberless" genome, not to be confused with a fiberless particle. Alternatively, a  
20 fiber protein may be encoded but is insufficiently expressed to result in a fiber containing particle.

Thus, among the delivery vectors contemplated for use are helper-independent fiberless recombinant adenovirus vector genomes that include genes that (a) express all adenovirus structural gene products but express insufficient  
25 adenovirus fiber protein to package a fiber-containing adenovirus particle without complementation of said fiber gene, (b) express an exogenous protein, and (c) contains an adenovirus packaging signal and inverted terminal repeats containing adenovirus origin of replication.

The adenovirus vector genome is propagated in the laboratory in the form  
30 of rDNA plasmids containing the genome, and upon introduction into an appropriate host, the viral genetic elements provide for viral genome replication

-54-

and packaging rather than plasmid-based propagation. Exemplary methods for preparing an Ad-vector genome are described in the Examples.

A vector herein includes a nucleic acid (preferably DNA) molecule capable of autonomous replication in a cell and to which a DNA segment, *e.g.*, a gene or polynucleotide, can be operatively linked to bring about replication of the attached segment. For purposes herein, one of the nucleotide segments to be operatively linked to vector sequences encodes at least a portion of a therapeutic nucleic acid molecule. As noted above, therapeutic nucleic acid molecules include those encoding proteins and also those that encode regulatory factors that can lead to expression or inhibition or alteration of expression of a gene product in a targeted cell.

#### 1. Nucleic Acid Gene Expression Cassettes

In various embodiments, a peptide-coding sequence of the therapeutic gene is inserted into an expression vector and expressed; however, it is also feasible to construct an expression vector which also includes some non-coding sequences as well. Preferably, however, non-coding sequences are excluded. Alternatively, a nucleotide sequence for a soluble form of a polypeptide may be utilized. Another preferred therapeutic viral vector includes a nucleotide sequence encoding at least a portion of a therapeutic nucleotide sequence operatively linked to the expression vector for expression of the coding sequence in the therapeutic nucleotide sequence.

The choice of viral vector into which a therapeutic nucleic acid molecule is operatively linked depends directly, as is well known in the art, on the functional properties desired, *e.g.*, vector replication and protein expression, and the host cell to be transformed. Although certain adenovirus serotypes are recited herein in the form of specific examples, it should be understood that the use of *any* adenovirus serotype, including hybrids and derivatives thereof are contemplated.

A translatable nucleotide sequence is a linear series of nucleotides that provide an uninterrupted series of at least 8 codons that encode a polypeptide in one reading frame. Preferably, the nucleotide sequence is a DNA sequence. The vector itself may be of any suitable type, such as a viral vector (RNA or DNA),

-55-

naked straight-chain or circular DNA, or a vesicle or envelope containing the nucleic acid material and any polypeptides that are to be inserted into the cell.

## 2. Promoters

As noted elsewhere herein, an expression nucleic acid in an Ad-derived  
5 vector may also include a promoter, particularly a tissue or cell specific promoter, preferably one expressed in the targeted cells.

Promoters nucleic acid fragments that contain a DNA sequence that controls the expression of a gene located 3' or downstream of the promoter. The promoter is the DNA sequence to which RNA polymerase specifically binds  
10 and initiates RNA synthesis (transcription) of that gene, typically located 3' of the promoter. A promoter also includes DNA sequences which direct the initiation of transcription, including those to which RNA polymerase specifically binds. If more than one nucleic acid sequence encoding a particular polypeptide or protein is included in a therapeutic viral vector or nucleotide sequence, more  
15 than one promoter or enhancer element may be included, particularly if that would enhance efficiency of expression. Regulatable (inducible) as well as constitutive promoters may be used, either on separate vectors or on the same vector.

For example, some useful regulatable promoters are those of the CREB-  
20 regulated gene family and include inhibin, gonadotropin, cytochrome c, glucagon, and the like. (See, e.g., International PCT application No. WO 96/14061). Preferably the promoter selected is from a photoreceptor-specific gene, such as a rhodopsin gene or gene that encodes a protein that regulates rhodopsin expression.

25 A regulatable or inducible promoter may be described as a promoter wherein the rate of RNA polymerase binding and initiation is modulated by external stimuli. (see, e.g., U.S. Patent Nos. 5,750,396 and 5,998,205). Such stimuli include various compounds or compositions, light, heat, stress, chemical energy sources, and the like. Inducible, suppressible and repressible promoters  
30 are considered regulatable promoters.

Regulatable promoters may also include tissue-specific promoters. Tissue-specific promoters direct the expression of the gene to which they are

-56-

operably linked to a specific cell type. Tissue-specific promoters cause the gene located 3' of it to be expressed predominantly, if not exclusively, in the specific cells where the promoter expressed its endogenous gene. Typically, it appears that if a tissue-specific promoter expresses the gene located 3' of it at all, then  
5 it is expressed appropriately in the correct cell types (see, e.g., Palmiter et al. (1986) Ann. Rev. Genet. 20: 465-499).

E. Formulation and administration

Compositions containing therapeutically effective concentrations of recombinant adenovirus delivery vectors for delivery of therapeutic gene  
10 products to cells that express the targeted receptor, such as IGF-1 receptors and TNF- $\alpha$  receptors.

Preferable modes of administration include, local and topical modes of administration, such as, but are not limited to, intramuscular, intravenous, intraperitoneal and subretinal injection, particularly intravitreal injection,

15 The recombinant viral compositions may also be formulated for implantation into tissues, including as the anterior or posterior chamber of the eye, particularly the vitreous cavity, in sustained released formulations, such as adsorbed to biodegradable supports, including collagen sponges, or in liposomes. Sustained release formulations may be formulated for multiple dosage  
20 administration, so that during a selected period of time, such as a month or up to about a year, several dosages are administered. Thus, for example, liposomes may be prepared such that a total of about two to up to about five or more times the single dosage is administered in one injection. The vectors are formulated in an pharmaceutically acceptable carriers for the selected route of  
25 administration in a volume suitable for such route.

To prepare compositions the viral particles are dialyzed into a suitable pharmaceutically acceptable carrier or viral particles may be concentrated and/or mixed therewith. The resulting mixture may be a solution, suspension or emulsion. In addition, the viral particles may be formulated as the sole  
30 pharmaceutically active ingredient in the composition or may be combined with other active agents for the particular disorder treated.

-57-

Suitable carriers include, but are not limited to, physiological saline, phosphate buffered saline (PBS), balanced salt solution (BSS), lactate Ringers solution, and solutions containing thickening and solubilizing agents, such as glucose, polyethylene glycol, and polypropylene glycol and mixtures thereof.

- 5 Liposomal suspensions may also be suitable as pharmaceutically acceptable carriers. These may be prepared according to methods known to those skilled in the art.

The compositions can be prepared with carriers that protect them against rapid elimination from the body, such as time release formulations or coatings.

- 10 Such carriers include controlled release formulations, such as, but not limited to, microencapsulated delivery systems, and biodegradable, biocompatible polymers, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, polyorthoesters, polylactic acid and other types of implants that may be placed directly into the body or tissue of interest.

- 15 Liposomal suspensions may also be suitable as pharmaceutically acceptable carriers. Preferably such liposomes. For example, liposome formulations may be prepared by methods known to those of skill in the art [see, e.g., Kimm *et al.* (1983) *Bioch. Bioph. Acta* 728:339-398; Assil *et al.* (1987) *Arch Ophthalmol.* 105:400; and U.S. Patent No. 4,522,811]. The viral particles  
20 may be encapsulated into the aqueous phase of liposome systems. The active materials can also be mixed with other active materials, that do not impair the desired action, or with materials that supplement the desired action or have other action, such as anti-tumor agents.

- For purposes herein, the viral or other particles may be complexed with  
25 the bifunctional molecules (the conjugates) prior to packaging or immediately prior to use. Hence combinations and kits containing the combinations of the selected delivery vector and the bifunctional molecules are also provided. The bifunctional molecules and delivery vectors may be packaged as separate compositions or as a single composition. The kits optionally include instructions  
30 for use and administration of the combinations.

The compositions can be enclosed in ampules, disposable syringes or multiple or single dose vials made of glass, plastic or other suitable material.

-58-

Such enclosed compositions can be provided in kits. In particular, kits containing vials, ampules or other container, preferably disposable vials or containers or other packages with sufficient amount of the composition to deliver a desired amount, which depends upon the treated condition.

5           Finally, the combinations and components compositions thereof may be packaged as articles of manufacture containing packaging material, typically a vial or container, an pharmaceutically acceptable composition containing the viral particles and a label that indicates the therapeutic use of the composition.

          Also provided are kits for practice of the methods herein. The kits  
10       contain one or more containers, such as sealed vials, with sufficient composition for single dosage administration.

#### Administration

          The compositions containing the compounds are administered systemically by any suitable route, or may be administered topically, such as by  
15       injection into synovial fluids for treatment of rheumatoid arthritis, in the form of penetrating eyedrops for treatment of ocular disorders or disorders in which the vectors can be suitably targeted from the eye.

          It is further understood that, for any particular subject and disorder, specific dosage regimens should be adjusted over time according to the  
20       individual need and the professional judgment of the person administering or supervising the administration of the recombinant viruses, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the methods provided herein.

#### F. Diseases, Disorders and therapeutic products

25           The bifunctional molecules provided herein permit targeting of viral and bacterial vectors to cells that express targeted receptors. The targeted receptors are those that activate the PI3K signalling pathway and internalized linked ligands by virtue thereof. Because such receptors are diverse and widespread, the use of bifunctional molecules provides a flexible means for gene delivery and  
30       therapeutic product delivery to cells and tissues. Receptors for the targeting agents, such as growth factors and hormones, are overexpressed on cells

-59-

associated with various disorders and conditions and/or on only selected cell types.

The use of the PI3K signaling strategy to tailor delivery of the adenovirus vectors to cell types over-expressing the desired growth factor receptor or cytokine receptor or other cell surface receptor that activates the PI3K pathway permits internalization via such receptors. Adenovirus binding and internalization can be further enhanced by taking advantage of additional interactions with CAR and also with integrins, if they are present on the cell surface. The bifunctional molecules provided herein do not bind the viral fiber protein, thus allowing for interactions with CAR. In addition, preferred bifunctional molecules bind the penton base protein, which has five "RGD" integrin binding sites. Since not all five "RGD" sites are bound by the whole molecule, the presence of the bifunctional molecule precludes neither CAR nor integrin interactions, and in addition offers a specific growth factor/cytokine - receptor binding interaction. In that sense, it is like a "triple-edged sword" for enhanced levels of binding and internalization, once targeting has been achieved. It is also advantageous to adopt a targeting strategy, which does not preclude interaction of the adenovirus with the  $\alpha v$  integrins because the integrins are known to not only signal on their own, but to promote optimal activation of growth factor receptors (Vuori *et al.*, *Science*, 266:1576 (1994); Cybulsky *et al.*, *J. Clin. Invest.*, 94:68 (1994); Jones *et al.*, *J. Cell Biol.*, 139:279 (1997); Miyamoto *et al.*, *J. Cell Biol.*, 135:1633 (1996); Schneller *et al.*, *EMBO J.*, 16:5600 (1997); Woodard *et al.*, *J. Cell Sci.*, 111:469 (1998); Moro *et al.*, *EMBO J.*, 17:6622 (1998); Soldi *et al.*, *EMBO J.*, 18:882 (1999)).

#### 25 1. Diseases and disorders

Methods for specifically targeting recombinant adenovirus vectors for delivery of gene products, particularly therapeutic products are provided herein. These methods are particularly suitable for targeting cells that express receptors to which the bifunctional molecules provided herein selectively bind resulting in internalization of the linked delivery vector. Adenoviruses are presently preferred. Diseases that can be targeted include, but are not limited to, cancers, vascular disorders, diabetic retinopathies, restenosis, ophthalmic disorders,



-60-

hyperproliferative disorders and hormonal disorders. The methods and bifunctional molecules provided herein permit targeting to restricted sets of cells associated with particular disorders.

#### Angiogenesis

5           In the normal adult, angiogenesis is tightly regulated and limited to wound healing, pregnancy and uterine cycling. Angiogenesis is turned on by specific angiogenic molecules such as basic and acidic fibroblast growth factor (FGF), vascular endothelial growth factor (VEGF), angiogenin, transforming growth factor (TGF), tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and platelet derived growth factor  
10 (PDGF). Angiogenesis can be suppressed by inhibitory molecules such as interferon- $\alpha$ , thrombospondin-1, angiostatin and endostatin. It is the balance of these naturally occurring stimulators and inhibitors that controls the normally quiescent capillary vasculature. When this balance is upset, as in certain disease states, capillary endothelial cells are induced to proliferate, migrate and  
15 ultimately differentiate.

Angiogenesis plays a central role in a variety of disease including cancer and neovascularization. Sustained growth and metastasis of a variety of tumors has also been shown to be dependent on the growth of new host blood vessels into the tumor in response to tumor derived angiogenic factors. Proliferation of  
20 new blood vessels in response to a variety of stimuli occurs as the dominant finding in the majority of eye disease and that blind including proliferative diabetic retinopathy (PDR), age-related macular degeneration (ARMD), rubeotic glaucoma, interstitial keratitis and retinopathy of prematurity. In these diseases, tissue damage can stimulate release of angiogenic factors resulting in capillary  
25 proliferation. VEGF plays a dominant role in iris neovascularization and neovascular retinopathies. While reports clearly show a correlation between intraocular VEGF levels and ischemic retinopathic ocular neovascularization, FGF likely plays a role. Basic and acidic FGF are known to be present in the normal adult retina, even though detectable levels are not consistently correlated with  
30 neovascularization. This may be largely due to the fact that FGF binds very tightly to charged components of the extracellular matrix and may not be readily

-61-

available in a freely diffusible form that would be detected by standard assays of intraocular fluids.

A final common pathway in the angiogenic response involves integrin-mediated information exchange between a proliferating vascular endothelial cell and the extracellular matrix. This class of adhesion receptors, called integrins, are expressed as heterodimers having an  $\alpha$  and  $\beta$  subunit on all cells. One such integrin,  $\alpha_v\beta_3$ , is the most promiscuous member of this family and allows endothelial cells to interact with a wide variety of extracellular matrix components. Peptide and antibody antagonists of this integrin inhibit angiogenesis by selectively inducing apoptosis of the proliferating vascular endothelial cells. Two cytokine-dependent pathways of angiogenesis exist and may be defined by their dependency on distinct vascular cell integrins,  $\alpha_v\beta_3$  and  $\alpha_v\beta_5$ . Specifically, basic FGF- and VEGF-induced angiogenesis depend on integrin  $\alpha_v\beta_3$  and  $\alpha_v\beta_5$ , respectively, since antibody antagonists of each integrin selectively block one of these angiogenic pathways in the rabbit corneal and chick chorioallantoic membrane (CAM) models. Peptide antagonists that block all  $\alpha_v$  integrins inhibit FGF- and VEGF-stimulated angiogenesis. While normal human ocular blood vessels do not display either integrin,  $\alpha_v\beta_3$  and  $\alpha_v\beta_5$  integrins are selectively displayed on blood vessels in tissues from patients with active neovascular eye disease. While only  $\alpha_v\beta_3$  was consistently observed in tissue from patients with ARMD,  $\alpha_v\beta_3$  and  $\alpha_v\beta_5$  were present in tissues from patients with PDR. Systemically administered peptide antagonists of integrins blocked new blood vessel formation in a mouse model of retinal vasculogenesis.

In addition to adhesion events described above, cell migration through the extracellular matrix also depends on proteolysis. Matrix metalloproteinases are a family of zinc-requiring matrix-degrading enzymes that include the collagenases, gelatinases and stromelysins, all of which have been implicated in invasive cell behavior. Invasive cell processes such as tumor metastasis and angiogenesis have been found to be associated with the expression of integrins and MMP-2, MMP-2 are all found throughout the eye where they may interact to maintain a quiescent vasculature until the balance is upset, resulting in pathological angiogenesis. A non-catalytic C-terminal hemopexin-like domain of MMP-2

-62-

(PEX) can block cell surface collagenolytic activity and inhibit angiogenesis in the CAM model by preventing localization of MMP-2 to the surface of invasive cells through interaction with the integrin  $\alpha_v\beta_3$ .

- Anti-angiogenic agents have a role in treating retinal degeneration to
- 5 prevent the damaging effects of trophic and growth factors. Angiogenic agents, also have role in promoting desirable vascularization to retard retinal degeneration by enhancing blood flow to cells.

#### Growth Factors/Cytokines and Pathological Conditions

##### EGF Receptors

- 10 EGF receptors are overexpressed in glioblastomas, bladder tumors, advanced gastric tumors and cervical cancers (Gullick (1991) *Br. Med. Bull.*, 47:87). They are overexpressed in 63% of tumor specimens from patients with lung cancer (Pastorino *et al.* (1993) *J. Cell. Biochem. Suppl.*, 17F:237), and are overexpressed in astrocytic gliomas (Goussia *et al.* (2000) *Oncol. Rep.* 7:401.
- 15 Overexpression is also correlated with tumor invasion and progression of human esophageal and gastric carcinomas (Yoshida *et al.*, *Exp. Pathol.*, 40:291 (1990)).

##### FGF Receptors

- These receptors are overexpressed in human pancreatic adenocarcinomas (Kobrin *et al.*, *Cancer Res.*, 53:4741 (1993)),
- 20 in human breast and gynecological cancers (Jaakkola *et al.*, *Int. J. Cancer*, 54:378 (1993)), human astrocytomas (Morrison *et al.*, *J. Neuro-oncol.*, 18:207 (1994)); and in human melanoma tissues (Xerri *et al.*, *Melanoma Res.*, 6:223 (1996)). FGF receptor expressing cells are also implicated in restenosis, Kaposi sarcoma, diabetic retinopathies and numerous disorders of the eye.

- 25 EGFR, FGFR and IGF-1R

EGFR, FGFR and IGF-1R are overexpressed in pancreatic cancers (Korc, *Surg. Oncol. Clin. N. Am.*, 7:25 (1998))

##### IGF-1R

- IFG-1R is overexpressed and hyperphosphorylated in primary breast
- 30 tumors (Surmacz, *J. Mamm. Gland Biol. Neoplasia*, 5:95 (2000)).

##### TNFR

-63-

Overexpression of both TNF $\alpha$  receptors, p55 and p75, is observed in neoplastic cells from patients with chronic lymphocytic leukemia (Waage *et al.*, *Blood*, 80:2577 (1992)) TNF $\alpha$  p55 receptor is overexpressed in breast carcinomas (Pusztai *et al.*, *Br. J. Cancer*, 70:289 (1994)). TNFR is overexpressed in normal  
5 and malignant myeloid cells, e.g., HL-60 promyelocytes (Munker *et al.*, *Blood*, 70:1730 (1987)).

#### SCF receptors

Development of mastocytosis (abnormal infiltration of mast cells into various organs) in patients with myelodysplastic syndrome (MDS) is due to a  
10 mutation in c-kit (the SCF receptor), which renders it ultra-sensitive to SCF (Dror *et al.*, *Br. J. Haematol.*, 108:729 (2000)). Therefore, SCF-derived bifunctional molecules can be used to deliver therapeutic products to such cells. Benign and malignant ovarian tumors express c-kit (SCF receptor) while normal tissue does not (Tonary *et al.*, *Int. J. Cancer*, 89:242 (2000)).

#### 15 2. Therapeutic products

Among the DNA that encodes therapeutic products contemplated for use is DNA encoding correct copies of defective genes, such as the defective gene (CFTR) associated with cystic fibrosis (see, e.g., International Application WO 93/03709, which is based on U.S. Application Serial No. 07/745,900; and  
20 Riordan *et al.* (1989) *Science* 245:1066-1073), and anticancer agents, such as tumor necrosis factors, and cytotoxic agents. Therapeutic products include but are not limited to, wild-type genes that are defective in targeted ocular disorders, such as defective gene products or fragments thereof sufficient to correct the genetic defect, trophic factors, including growth factors, inhibitors and agonists  
25 of trophic factors, anti-apoptosis factors and other products herein or known to those of skill in the art to be useful for treatment of selected disorders.

The following examples are included for illustrative purposes only and are not intended to limit the scope of the invention.

-64-

## EXAMPLE 1

## Preparation and characterization of DAV-1

A monoclonal antibody, designated DAV-1, had been previously obtained and characterized (see, Stewart *et al.* (1997) *EMBO J.* 16:1189-1198) as described in this Example. The nucleic acid and protein sequences of the heavy and light chain of DAV-1 are set forth in SEQ ID Nos. 1-4; and the nucleic acid and protein sequences of the portion used in exemplified fusion proteins containing substantially full-length heavy chain is set forth in SEQ ID Nos. 5 and 6.

## 10 Materials and Methods

## Cell lines, viruses and recombinant proteins

A549, HeLa, H2981 and SW480 cell lines, adenovirus serotypes Ad2, Ad3 and Ad4 were purchased from the American Type Tissue Culture Collection (Rockville, MD). For virus isolation, HeLa cells were infected with either Ad2, Ad3 or Ad4 at a multiplicity of 10 p.f.u./cell and then harvested 48-72 h later. Cells were frozen and thawed five times to release intracellular virus particles. After removing the cell debris by high-speed centrifugation, virions were isolated by banding on a 15-40% cesium chloride gradient in 10 mM Tris-HCl, 150 mM NaCl pH 8.1 (TBS) as reported previously (Everitt *et al.*, 1977). Banded virions were removed and then dialyzed into TBS buffer containing 10% glycerol, except for cryo-EM studies in which case the CsCl was removed by multiple centrifugation steps in a Microcon 100 (Amicon) filtration device using pH 8.1 phosphate buffer. Recombinant Ad2 penton base containing 571 amino acids (Neumann *et al.*, 1988) was produced in Sf9 insect cells using baculovirus as previously described (Nemerow *et al.*, 1993); Wickham *et al.*, 1993).

## 25 Generation and characterization of the DAV-1 anti-penton base mAb

A hybridoma (designated DAV-1) secreting a mAb of the subtype  $\gamma 1\kappa$  was generated by standard techniques. The DAV-1 mAb was purified from ascites fluids using protein G-Sepharose (HiTrap GII, Pharmacia). Fab fragments of the DAV-1 mAb were generated by papain digestion. Briefly, 1-5 mg/ml of purified DAV-1 IgG in 50 mM Tris-HCl pH 8.0, 10 mM L-cysteine, 3 mM EDTA was incubated for 7 h at 37°C in the presence of 8% w/w soluble papain

-65-

(Sigma Chemical Co., St. Louis, MO). The reaction was stopped by the addition of 30 mM iodoacetamide, and the Fab antibody fragments were then isolated on a Resource S FPLC column (Pharmacia) equilibrated with 50 mM MES pH 5.0. The purified Fab fragments were analyzed by SDS-PAGE and then concentrated  
5 to 2.2 mg/ml using a Centricon 10 membrane ultrafiltration device (Amicon).

Reactivity of the DAV-1 mAb with different adenovirus serotypes was quantified in an ELISA. Ninety-six well polystyrene plates (Immobilon, Dynatech) were coated with 1  $\mu$ g of penton base or with 5  $\mu$ g of purified Ad2, Ad3 or Ad4 in PBS for 18 h at 4°C. After blocking non-specific binding sites with 2% non-  
10 fat dried milk, 10  $\mu$ g/ml of purified DAV-1 mAb or an irrelevant control antibody were added to the wells for 60 min at 22°C. Antibody binding was detected by the addition of alkaline phosphatase linked to goat anti-mouse IgG followed by substrate (Sigma Chemical Co., St. Louis, MO). Substrate development was quantified at 405 nm in an ELISA plate reader (Titertek, Flow laboratories).

15 To examine whether the DAV-1 mAb also recognized RGD-containing cell matrix proteins, 1-2  $\mu$ g of recombinant penton base protein, or fibronectin, vitronectin, collagen (type 1) and fibrogen were reacted with the DAV-1 mAb in a Western blot. Recombinant penton base or cell matrix proteins were electrophoresed on a 8-15% gradient SDS gel (Novex, San Diego, CA) and then  
20 transferred to a nitrocellulose filter (Immobilon P, Millipore). Following blocking of non-specific binding sites with 1% non-fat dried milk (Blotto), the filters were reacted with 10  $\mu$ g/ml of the DAV-1 mAb followed by incubation with alkaline phosphatase linked to goat anti-mouse IgG (Tropix, Bedford, MA) and then with a chemiluminescent substrate (CDP).

#### 25 Functional analysis of the DAV-1 mAb

The effect of the DAV-1 mAb on penton base binding to cell surface  $\alpha_v$  integrins was examined as follows. To  $1 \times 10^6$  A549 epithelial cells 10  $\mu$ g/ml of purified DAV-1 IgG or Fab fragments of the DAV-1 mAb was added. Varying amounts of  $^{125}$ I-labeled penton base (10  $\mu$ Ci/ $\mu$ g) were then added to the cells in  
30 the presence or absence of a 50-fold excess of unlabeled penton base and incubated for 60 min at 4°C. Unbound penton base was removed by centrifuging the cell samples through a cushion of 1:1 glycerol/mineral oil and

-66-

the amount of cell-associated penton base was determined by counting the cell pellet in a  $\gamma$ -counter.

The effect of the DAV-1 mAb on adenovirus infection was quantified by plaque assay. A549 cells were seeded into six-well plates and cultured to 90% confluency. Three  $\mu\text{g/ml}$  of DAV-1 Fab or 18  $\mu\text{g}$  of whole IgG DAV-1 antibody were added to the cell cultures, followed by addition of purified 100 p.f.u. Ad2 and incubation at 37°C for 2 h. The Ad2 and antibody mixtures were then removed, and 8 ml of overlay medium containing 0.5% agarose in DMEM medium and 10% FCS was added into each well. The cells were fed with 4 ml of overlay medium on day 5 post-infection. The plaques were scored on day 10 post-infection.

#### Epitope mapping and kinetic analysis of DAV-1 binding to penton base

As noted above, the DAV-1 binding site on the Ad2 penton base was identified by affinity-directed mass spectrometry. For these studies, a region of the penton base that approximately spanned the RGD-containing epitope sequence was selected. A series of overlapping synthetic peptides varying by one amino acid on the N-terminal or C-terminal region of the Ad2 penton base RGD sequence,  $^{480}\text{MNDHAIRGDTFATRA}^{494}$  (SEQ ID NO. 19), was generated by solid-phase protocols, and the precise boundaries of the DAV-1 epitope were then determined by affinity-directed mass spectrometry (see, Zhao *et al.* (1994) *Anal. Chem.* 66:3723-3726; Zhao *et al.* (1996) *Proc. Natl. Acad. Sci. U.S.A.* 93:4020-4024).

Precise measurements of DAV-1 interactions with the penton base protein were determined by SPR (Karlsson *et al.* (1991) *J. Immunol. Methods* 145:229-240) using an automated biosensor system (BIAcore 2000 Pharmacia). Briefly, recombinant penton base at 60  $\mu\text{g/ml}$  was immobilized onto carboxymethyl dextran-coated biosensor chips in 10 mM MES pH 6.5 containing 10 mM NaCl. Following amine coupling of the penton base, varying amounts of purified Fab fragments (3.6-57.0  $\mu\text{g/ml}$ ) or IgG molecules (36-576  $\mu\text{g/ml}$ ) were flowed over the penton base at a rate of 40  $\mu\text{l/min}$ , respectively. Kinetic binding data ( $K_{\text{on}}$ ,  $K_{\text{off}}$  and  $K_D$ ) were obtained using BIAevaluation software (version 2.1). Stoichiometric data was obtained by observing the change in SPR at saturation

-67-

binding and assuming a molecular mass of 350 kDa for the penton base, 43 kDa for Fab fragments and 150 kDa for IgG molecules. The sequence recognized by DAV-1 was IRGDTFATR (see SEQ ID NO. 20).

## EXAMPLE 2

### 5 Preparation and Analysis of Bifunctional Signaling Antibodies

#### A. Cloning of DAV-1-encoding cDNA

The hybridoma secreting a monoclonal antibody of the type  $\gamma 1\kappa$  (designated DAV-1) was generated as described in EXAMPLE 1 (see, Stewart *et al.* (1997) *EMBO J.* 16:1189-1198). Total RNA was isolated from the DAV-1 hybridoma cell line (Trizol reagent, Gibco BRL) and cDNA was generated using the Superscript Plasmid System kit (Gibco BRL) essentially as described (Uematsu, *Immunogenetics*, 34: 174-178 (1991)). The DAV-1 heavy chain was PCR amplified with: primers 5'-CCT GCT CTG TGT TTA CAT GAG GG (CH3 region) (SEQ ID. NO. 15); 5'-CCC AGG GTC ATG GAG TTA G (CH1 region) (SEQ ID. NO. 16); for kappa (light) chain amplification: 5'-AAG ATG GAT ACA GTT GGT GC (CL-A) (SEQ ID. NO. 17) and 5'-TGT CAA GAG CTT CAA CAG GA (CL-B) (SEQ ID. NO. 18) were the primers used. PCR amplification was carried out using the parameters 94°C for 1 minute, 63°C for 1 minute and 72°C for 1 minute, for a total of 30 cycles. The PCR products were ligated into pCR2.1 (Invitrogen, Carlsbad, CA) using a TA cloning kit (Invitrogen, Carlsbad, CA), and then sequenced by automated sequencing. Further amplification to obtain the complete DAV-1 heavy and light chains was performed using standard PCR reactions as described above.

25 The cDNA coding the  $\gamma$  heavy and  $\kappa$ -light chains of (DAV-1) were subcloned into an expression vector, designated pIZ, which contains a Zeocin selection marker for expression in insect cells (Invitrogen, CA). The portion of the DAV-1 heavy chain used to generate fusion proteins with various cytokines or growth factors was the full-length heavy chain minus the last 18 amino acids (coding portion in SEQ ID NO. 1 encoding amino acid 1 through 438, and SEQ ID NO. 5) was cloned into the pIZ expression vector between its KpnI and EcoRI restriction sites by introducing KpnI and EcoRI sites into the 5' and 3' ends, respectively. (see, SEQ ID. NO. 5). This was accomplished by using the "sense"



-68-

PCR primer 5'-GGT ACC GCC ACC ATG GGA TGG AGC TGG ATC T (SEQ ID. NO. 21) having a KpnI restriction site, and the "antisense" primer 5'-GAA TTC ATG TAA CAC AGA GCA GGA (SEQ ID. NO. 22) having an EcoRI restriction site, for PCR amplification of the portion of the DAV-1 heavy chain sequence set forth in SEQ ID. NO. 5, prior to cloning. The DAV-1 light chain (SEQ ID NO. 3) was cloned into the HindIII and XbaI sites of the pIZ expression vector by introducing HindIII and XbaI sites into the 5' and 3' ends of the light chain encoding DNA (see SEQ ID. NO. 3). This was accomplished by using the "sense" PCR primer 5'-AAG CTT GCC ACC ATG GAG ACA GAC ACA ATC CTG CT (SEQ ID. NO. 23) having a HindIII restriction site, and the "antisense" primer 5'-TCT AGA TGT CTC TAA CAC TCA TTC CTG T (SEQ ID. NO. 24) having an XbaI restriction site, for PCR amplification of the DAV-1 light chain sequence set forth SEQ ID. NO. 3, prior to cloning. The resulting vectors were designated as pIZ- $\gamma$ , pIZ- $\kappa$  for heavy chain and light chain constructs, respectively.

Fab'<sub>2</sub> forms of the DAV-1 antibody were also PCR amplified by creating a C-terminal deletion in the DAV-1  $\gamma$ -heavy chain at amino acid position 247 (see, SEQ ID No. 1), and were cloned between the KpnI and EcoRI sites of the pIZ vector using the "sense" primer sequence set forth in SEQ ID. NO. 21 (having a KpnI restriction site), and the "antisense" primer 5'-GAA TTC TGA TAC TTC TGG GAC TGT (SEQ ID. NO. 25 with an an EcoRI restriction site).

#### B. Cloning of the Bifunctional Signaling Antibodies

DNA encoding full length mature human TNF- $\alpha$ , IGF-1, EGF and SCF peptides were obtained by PCR amplification of: cDNA obtained from ATCC (Rockville, Md) encoding human TNF- $\alpha$ ; RT-PCR of total RNA isolated from U937 cells for human IGF-1; cDNA obtained from Invitrogen, Carlsbad, CA for SCF; and a synthetic template prepared by annealing two oligonucleotides having an 18 bp overlap (SEQ ID NOS. 30 and 31) for EGF. For construction of the fusion proteins, DNA sequences encoding the full length mature human TNF- $\alpha$ , IGF-1 and EGF peptides (peptide sequences set forth in SEQ ID. NOS. 7, 8 and 9 respectively) were ligated in frame into the EcoRI site in pIZ- $\gamma$  downstream (3'-end) of the DAV-1 heavy chain portion (SEQ ID. NO. 5) used to prepare the fusion proteins.

-69-

To facilitate cloning of the full length mature peptides into the pIZ- $\gamma$  vector at the EcoRI site 3' to the DAV-1 heavy chain sequence, EcoRI sites were introduced at the 5' and 3'-ends of the nucleic acid encoding each of the full length mature peptides TNF- $\alpha$ , IGF-1 and EGF by PCR amplification using the

5 following primers: For amplification of TNF- $\alpha$ , "sense" primer 5'-GAA TTC GTC AGA TCA TCT TCT CGA AC (SEQ ID. NO. 26) and "antisense" primer 5'-GAA TTC TAC AGG GCA ATG ATC CCA AA (SEQ ID. NO. 27); for amplification of IGF-1, "sense" primer 5'-GAA TTC GGA CCG GAG ACG CTC TGC GG (SEQ ID. NO. 28) and "antisense" primer 5'-GAA TTC TAA GCT GAC TTG GCA GGC TT

10 (SEQ ID. NO. 29); for amplification of EGF, "sense" primer 5'-GAA TTC AAT AGT GAC TCT GAA TGT CCC CTG TCC CAC GAT GGG TAC TGC CTC CAT GAT GGT GTG TGC

ATG TAT ATT GAA GCA TTG GAC AAG TAT GCA (SEQ ID. NO. 30) and "antisense" primer 5'-GAA TTC TAG CGC AGT TCC CAC CAC TTC AGG TCT

15 CGG TAC TGA CAT CGC TCC CCG ATG TAG CCA ACA ACA CAG TTG CAT GCA TAC TTG TCC AAT GCT TC (SEQ ID. NO. 31). The orientation of the fusion proteins was determined by PCR analysis. All sequences were confirmed using automated DNA sequencing.

The fusion proteins contain the portion of the DAV-1 heavy chain

20 sequence set forth in SEQ ID. NO. 6, followed by a two amino acid "linker" sequence generated by the EcoRI site between the DAV-1 and growth factor sequences (see amino acids 439 (Glu) and 440 (Phe) in SEQ ID NOS. 11, 12 and 13), followed in-frame by the full length, mature growth factor peptide. Sequences of fusion proteins of DAV-1 heavy chain with TNF- $\alpha$ , IGF-1 and EGF

25 are set forth in SEQ ID. NOS. 11, 12 and 13, respectively.

For construction of the fusion protein of DAV-1 heavy chain with SCF, the DNA sequence encoding the full length mature SCF peptide (peptide sequence set forth in SEQ ID. NO. 10) was ligated in frame between the NotI and XbaI sites of the pIZ- $\gamma$  vector, downstream (3'-end) of the DAV-1 heavy

30 chain portion (SEQ ID. NO. 5) used to prepare the fusion protein. To facilitate cloning of the full length mature SCF peptide into the pIZ- $\gamma$  vector at the NotI and XbaI sites 3' to the DAV-1 heavy chain sequence, NotI and XbaI sites were

-70-

introduced at the 5' and 3'-ends of the SCF sequence encoding the full length mature peptide by PCR amplification using the following primers: "sense" primer 5'-GCG GCC GCA AGG GAT CTG CAG GAA TCG (SEQ ID. NO. 32) and "antisense" primer 5'-TCT AGA GTG CAA CAG GGG GTA ACA TA (SEQ ID.

5 NO. 33). Generation of the NotI site at the 5'-end of the full length SCF-encoding nucleic acid resulted in change of the first amino acid from glutamic acid (Glu) in the wildtype sequence (see first amino acid of SEQ ID. NO. 10) to glutamine (Gln) in the fusion protein (see amino acid 450 in SEQ ID. NO. 14). The resulting fusion construct includes the portion of the DAV-1 heavy chain  
10 peptide sequence set forth in SEQ ID. NO. 6, followed by 11 amino acids of "linker" sequence (amino acids 439 to 449 in SEQ ID. NO. 14), followed by the SCF mature peptide. The sequence of the resulting fusion protein is set forth in SEQ ID. NO. 14.

Similar cloning strategies were employed to generate growth factors such  
15 as TNF- $\alpha$ , IGF-1, EGF and SCF fused to the 3'-end of the Fab'<sub>2</sub> forms of the DAV-1 antibody.

#### C. Generation of secreted bifunctional antibody fusion proteins

The DAV-1 heavy chain-TNF, heavy chain-IGF-1 or heavy chain-EGF expression vector and the DAV-1 light chain expression vector were co-  
20 transfected into SF9 insect cells. The SF9 insect cells (Invitrogen) were transfected with a total of 3  $\mu$ g plasmid DNA comprising the heavy chain-growth factor / cytokine fusion and light chain vectors using 15  $\mu$ l Superfect (Qiagen) in 200  $\mu$ l DMEM (serum-free) at room temperature for 15 minutes and then added to fresh cultures of SF9 cell monolayers (about 90% confluency). Transfected  
25 cells were then subjected to Zeocin selection (600  $\mu$ g/ml) for the production of bifunctional molecules using a penton base ELISA assay. A pool of positive Zeocin-resistant cells was then selected for the production of fusion proteins.

#### D. Purification and functional analysis of bifunctional molecules

Supernatants from transfected SF9 cells were assayed for bifunctional  
30 antibody production by an ELISA assay using immobilized penton base as previously described (Mathias *et al.*, *J. Virol.*, 72(11): 8669 (1998)) and as described in this Example. Recombinant Ad2 penton base was produced in

-71-

- Trichoplusia Tn 5B1-4 insect cells (Wickham *et al.*, *Biotech. Prog.*, 8:391-396 (1992)) using the baculovirus vector pBlueBac (Invitrogen, Carlsbad, CA). The recombinant Ad2 penton base was used to coat a 96-well plastic tissue culture plate (Immulon-4, Dynatech) at a concentration of 1  $\mu$ g protein/well. Non-specific binding sites were quenched by incubation with a blocking agent (Superblock; Pierce), and then the SF9 cell culture supernatants were added to the plates and incubated at 22 °C for 1 h. Bound antibody was detected with HRP conjugated rabbit anti-mouse antibody (whole molecule). Culture supernatants that tested positive for the bifunctional antibody were passed through a Protein L (Actigen, Cambridge, UK) affinity column. Bound antibody was eluted with either 50 mM diethylamine (pH 9.7) or 0.1 M citrate-0.15 M NaCl (pH 3.0), and pooled fractions were dialyzed against PBS (10 mM sodium phosphate, 150 mM NaCl, pH 7.2). The purified fusion proteins were further characterized by SDS-PAGE and western blot.
- 15        The cytotoxic activity of the DT bifunctional monoclonal antibodies (DAV-1-TNF $\alpha$  bifunctional antibody) resulting from its interaction with TNF- $\alpha$  receptors was assayed using the TNF-sensitive MCF-7 cell line as previously described (Xiang *et al.*, *J. Biotech.*, 53: 3 (1997).
- 20        SF9 cell supernatants or the purified bifunctional proteins were separated on 12% SDS-PAGE gels and either stained with Coomassie blue or transferred to PVDF membrane filters (Amersham) and probed with rabbit anti-mouse (Sigma, St. Louis, MO) or goat anti-human TNF- $\alpha$  (Chemicon, Temecula, CA) antibodies followed by secondary antibodies conjugated to horseradish peroxidase (Sigma, St. Louis, MO) and detection with a chemiluminescence reagent (Supersignal, Pierce, Rockford, IL). Western blot analyses showed that the DAV-1 monoclonal antibody (D molecule) and the DAV-1-TNF $\alpha$  bifunctional antibody (DT molecule) were both recognized by an anti-mouse IgG antibody, while only the DT molecule was recognized by an anti-TNF $\alpha$  polyclonal antibody.
- 25

-72-

**EXAMPLE 3****Gene delivery vectors and complexing with bifunctional molecules****A. Adenovirus propagation**

Cell lines that are commonly used for growing adenovirus are useful as  
5 host cells for the preparation of adenovirus packaging cell lines. Preferred cells  
include 293 cells, an adenovirus-transformed human embryonic kidney cell line  
obtained from the ATCC, having Accession Number CRL 1573; HeLa, a human  
epithelial carcinoma cell line (ATCC Accession Number CCL-2); A549, a human  
lung carcinoma cell line (ATCC Accession Number CCL 1889); and other  
10 epithelial-derived cell lines. As a result of the adenovirus transformation, the  
293 cells contain the E1 early region regulatory gene. All cells were maintained  
in complete DMEM + 10% fetal calf serum unless otherwise noted.

These cell lines allow the production and propagation of adenovirus-based  
gene delivery vectors that have deletions in preselected gene regions and that  
15 are obtained by cellular complementation of adenoviral genes. Such units  
include but are not limited to E1 early region, E4 and the viral fiber gene.

Adenovirus type 2 (Ad2, ATCC) was propagated in A549 (ATCC # CCL  
185) epithelial cells and purified as previously described (Wickham *et al.*, *Cell*,  
73: 309 (1993)) and as described in this Example. Cells were infected with Ad2  
20 at a multiplicity of infection (MOI) of 10 and then harvested 2-3 days later. Cells  
were frozen and thawed five times to release intracellular particles, and then the  
cell debris was removed by centrifugation. The cell lysate was subjected to  
density gradient-ultracentrifugation on 25%-40% cesium chloride gradients, and  
the virus band was removed and dialyzed against 40 mM Tris-HCl-buffered  
25 saline, pH 8.1, containing 10% glycerol.

**B. Adenovirus gene delivery vectors**

Adenoviral vectors for delivery of genes, such as therapeutic genes, and  
methods for their construction and propagation are well known and readily  
available (see, *e.g.*, co-pending U.S. application Serial No. 09/482,682;  
30 International PCT application No. PCT/US00/00265, and U.S. application Serial  
No. 09/562,934). Other delivery vectors may be used. A variety of Ad delivery  
vectors are known and available.

-73-

In general, exemplary fiber-expressing and fiberless recombinant adenovirus vectors have been described (Von Seggern *et al.*, *J. Virol.*, 73: 1601 (1999); copending U.S. application Serial No. 09/482,682 filed January 14, 2000, and also International PCT application No. PCT/US00/00265, filed 5 January 14, 2000)). Construction of Ad5. $\beta$ gal.wt and Ad5. $\beta$ gal. $\Delta$ F (deposited on January 15, 1999, the ATCC under accession number VR2636) is described therein. Ad2 or Ad5 vectors using the RSV LTR in place of the SV40 promoter can be constructed in a manner analogous to the similar Ad5-based vectors described in the above-noted applications. Such vectors were used in the 10 experiments in the Examples herein. As described therein, gutted Ad vectors are those from which most or all viral genes have been deleted. They are grown by co-infection of the producing cells with a "helper" virus (using an E1-deleted Ad vector). The helper virus *trans*-complements the missing Ad functions, including production of the viral structural proteins needed for particle 15 assembly. The helper virus can be a fiber-deleted Ad (such as that described in Von Seggern *et al.*, *J. Virol.* 73:1601-1608 (1999)). The vector is prepared in a fiber expressing cell line (see, *e.g.*, Von Seggern *et al.* (1998) *J. Gen. Virol.* 79:1461-1468; Von Seggern *et al.* (2000), *J. Virol.* 74:354-362). All the necessary Ad proteins except fiber are provided by the fiber-deleted helper virus, 20 and the particles are equipped with the particular fiber expressed by the host cells.

A helper adenovirus vector genome and a gutless adenoviral vector genome are delivered to a packaging cell line (see, *e.g.*, International PCT application No. PCT/US00/00265, filed January 14, 2000). The cells are 25 maintained under standard cell maintenance or growth conditions, whereby the helper vector genome and the packaging cell together provide the complementing proteins for the packaging of the adenoviral vector particle. Such gutless adenoviral vector particles are recovered by standard techniques. The helper vector genome may be delivered in the form of a plasmid or similar 30 construct by standard transfection techniques, or it may be delivered through infection by a viral particle containing the genome. Such viral particle is

-74-

commonly called a helper virus. Similarly, the gutless adenoviral vector genome may be delivered to the cell by transfection or viral infection.

- The helper virus genome is preferably a fiberless adenovirus vector genome. Preferably, such genome also lacks the genes encoding the adenovirus
- 5 E1A and E1B proteins. More preferably, the genome further lacks the adenovirus genes encoding the adenovirus E3 proteins. Alternatively, the genes encoding such proteins may be present but mutated so that they do not encode functional E1A, E1B and E3 proteins. Furthermore, such vector genome may not encode other functional early proteins, such as E2A, E2B, and E4 proteins.
- 10 Alternatively, the genes encoding such other early proteins may be present but mutated so that they do not encode functional proteins.

- The packaging cell also provides proteins necessary for the complementation of the gutless vector so that an adenovirus particle containing the gutless vector genome may be produced. Thus, the packaging cell line can
- 15 provide wild-type or modified fiber protein. Alternatively, the cell line could package a fiberless particle, which could be used by itself or to which exogenously provided fiber could be added.

- In producing gutless vectors, the helper virus genome is also packaged, thereby producing helper virus. In order to minimize the amount of helper virus
- 20 produced and maximize the amount of gutless vector particles produced, it is preferable to delete or otherwise modify the packaging sequence in the helper virus genome, so that packaging of the genome is prevented or limited. Since the gutless vector genome will have a packaging sequence, it will be preferentially packaged.
- 25 One way to do this is to mutate the packaging sequence by deleting one or more of the nucleotides comprising the sequence or otherwise mutating the sequence to inactivate or hamper the packaging function. An alternative approach is to engineer the helper genome so that recombinase target sites flank the packaging sequence and to provide a recombinase in the packaging cell. The action of
- 30 recombinase on such sites results in the removal of the packaging sequence from the helper virus genome. Preferably, the recombinase is provided by a nucleotide sequence in the packaging cell that encodes the recombinase. Most

-75-

preferably, such sequence is stably integrated into the genome of the packaging cell. Various kinds of recombinase are known by those skilled in the art. The preferred recombinase is Cre recombinase, which operates on so-called lox sites, which are engineered on either side of the packaging sequence as discussed  
5 above. Further information about the use of Cre-loxP recombination is found in U.S. Pat. No. 5,919,676 and Morsy and Caskey, *Molecular Medicine Today*, Jan. 1999, pgs. 18-24.

As the gutless vectors lack many or all Ad genes, they must be grown as mixed cultures in the presence of a helper virus which can provide the missing  
10 functions. To date, such helper viruses have provided all Ad functions except E1, and E1 is complemented by growth in 293 cells or the equivalent. The resulting virus particles are harvested, and the helper virus is typically removed by CsCl gradient centrifugation (the vector chromosome is generally shorter than the helper chromosome, resulting in a difference in buoyant density between the  
15 two particles).

An example of a gutless gene delivery vector is pAd $\Delta$ RSVDys (Haecker *et al.* (1996) *Human Gene Therapy* 7:1907-1914). This plasmid contains a full-length human dystrophin cDNA driven by the RSV promoter and flanked by Ad inverted terminal repeats and packaging signals. Desired therapeutic proteins  
20 and other products intended for delivery to cells can be readily substituted for the dystrophin gene in this vector.

293 cells are infected with a first-generation Ad which serves as a helper virus, and then transfected with purified vector DNA. The helper Ad genome and the delivery vector DNA are replicated as Ad chromosomes, and packaged  
25 into particles using the viral proteins produced by the helper virus. Particles are isolated and the delivery vector-containing particles separated from the helper by virtue of their smaller genome size and therefore different density on CsCl gradients.

The vector is grown in either 633 or 705 cells and Ad5. $\beta$ gal. $\Delta$ F is used  
30 as a helper virus; both helper and the delivery vector genomes replicate and are packaged into particles. The provides all the essential Ad proteins except fiber, and the fiber protein is that produced by the cells (Ad5 fiber in 633 cells and



-76-

Ad37 fiber in the case of 705 cells). The packaged viral particles are then isolated by centrifugation.

For experiments exemplified herein, the following first generation recombinant virus and its fiberless derivative were used. The first-generation  
5 Ad2 virus Ad.RSV $\beta$ -gal is an E-1 and E-3 deleted, replication-defective, recombinant vector containing a Rous Sarcoma virus regulatory sequence - driven *LacZ* reporter gene (Stratford-Perricaudet *et al.*, *J. Clin. Invest.*, 90:626 (1992)).

First, a recombinant plasmid pAd.RSV $\beta$ -gal was constructed, in which the  
10 *LacZ* gene with the SV40 early region polyadenylation signal driven by the Rous Sarcoma virus long terminal repeat (RSV-LTR) is inserted downstream of the 1.3 map units (mu) from the left end of the adenovirus type 5 (Ad5) genome in place of E1a and E1b (mu 1.3-9.4). The reporter gene is followed by mu 9.4-17 of  
15 Ad5 to allow homologous recombination with the adenoviral genome for the generation of the recombinant adenovirus. The recombinant adenovirus was constructed by *in vivo* homologous recombination between plasmid pAd.RSV $\beta$ -gal and Ad5 dl327, an E-3 deletion mutant of Ad5 (Trousdale, M.D. *et al.*, *Cornea*, 14:280 (1995)).

Briefly, cells were cotransfected with 5  $\mu$ g of linearized pAd.RSV $\beta$ -gal and  
20 5  $\mu$ g of the 2.6 - 100 mu fragment of Ad5 DNA. After overlaying with agar and incubation for 10 days at 37°C, plaques containing recombinant adenovirus were picked and screened for nuclear  $\beta$ -galactosidase activity. The Ad.RSV $\beta$ -gal/ $\Delta$ F vector is identical to the first-generation virus with the exception of a fiber deletion to generate fiberless virions.

25 As noted, these and other exemplary fiber-expressing and fiberless recombinant adenovirus vectors have been described (Von Seggern *et al.*, *J. Virol.*, 73: 1601 (1999); copending U.S. application Serial No. 09/482,682 filed January 14, 2000, and also International PCT application No. PCT/US00/00265, filed January 14, 2000)).

-77-

C. Complexing of adenovirus delivery vector particles with bifunctional molecules

Complexation of the recombinant adenovirus vectors with the bifunctional molecules was accomplished as follows: The E1 deleted adenovirus vector encoding *LacZ*, under control of the RSV LTR, in place of E1 (Ad.RSV. $\beta$ -gal vector described above), was incubated with DAV-TNF (DT molecules) or control DAV (D molecules) antibodies at a ratio of 2 antibody molecules per RGD motif of adenovirus at room temperature for 30 minutes in either EXEL-400 medium (JRH Bioscience, Lenexa, KS) or Dulbecco modified Eagle's medium (DMEM) (Gibco-BRL).

EXAMPLE 4

Adenovirus, gene express, cell binding and internalization assays

A. Gene delivery and expression assay

The complex of Example 3C was then mixed with M21-L12 human melanoma cells, which are deficient in  $\alpha_v\beta_3$  and  $\alpha_v\beta_5$  integrin (Wickham *et al.*, *Cell*, 73: 309 (1993)), on ice for 60 min. Unbound virus was removed by washing with ice-cold PBS, and the cells were warmed to 37 °C for varying times and then plated in tissue culture plates. Ad-mediated gene transfer was examined 24 or 48 hours post infection by staining for  $\beta$ -galactosidase activity as previously described, by incubating the cells for 60 min at 37 °C with 3.5 mM *o*-nitrophenyl- $\beta$ -D-galactopyranoside (ONPG) as a chromagenic substrate and in buffer containing 0.5% Nonidet P-40 (Li *et al.*, *J. Virol.*, 72: 2055 (1998); Huang *et al.*, *J. Virol.*, 70: 4502 (1996)).

B. Binding assay

To measure adenovirus binding to cells, 500  $\mu$ g aliquots of adenovirus vector was labeled with  $^{125}$ I by incubation with Iodogen (Pierce) in an Iodogen-coated tube containing 1 mCi of Na $^{125}$ I. Iodinated proteins were separated from free  $^{125}$ I by gel filtration as described (Huang *et al.* (1999) *J. Virol.* 73:2798-2802). Binding of radiolabeled adenovirus on M21-L12 cells was then quantitated as described earlier (Huang *et al.* (1999) *J. Virol.* 73:2798-2802), and as described in this Example.  $10^6$  cells in suspension were incubated with  $10^6$  cpm (viral particles were generally labeled to a specific activity of  $5 \times 10^6$  to

-78-

8 x 10<sup>6</sup> cpm/ $\mu$ g) of the labeled virus proteins at 4 °C for 2 h. Non-specific binding was determined by incubating cells and labeled proteins in the presence of a 100-fold excess of unlabeled virus or fiber protein. After the cells were washed four times in ice-cold phosphate buffered saline, specific binding was calculated by subtracting the non-specific binding from the total bound cpm.

#### C. Internalization assay

To measure virus endocytosis, cells were incubated with complexed <sup>125</sup>I-labeled adenovirus and DAV-TNF or control antibodies, in DMEM supplemented with 0.5% purified bovine serum albumin (BSA) at 4 °C for 60 minutes. After removal of non-bound virus by washing with ice-cold PBS, the cells were warmed to 37 °C for varying times. Uninternalized virions were removed by incubation with trypsin-EDTA at room temperature for 5 min. prior to counting the cell pellets.

#### EXAMPLE 5

Adenovirus complexed with bifunctional molecules targeted to receptors that promote internalization of ligands by PI3K activation are internalized via binding to the targeted receptors.

##### Analysis of bifunctional signaling antibodies.

DAV-1 bifunctional signaling antibodies, designated DT (DAV-1 fused to TNF- $\alpha$ ), DI (DAV-1 fused to IGF-1), and DE (DAV-1 fused to EGF), were expressed in insect cells as secreted proteins and purified on Protein L affinity columns. The DT heavy chain (D<sub>H</sub>T) had an apparent molecular weight of approximately 70kDa, consistent with the combined sizes of the DAV-1  $\gamma$  heavy chain (53 kDa) and monomeric TNF- $\alpha$  ligand (17 kDa). The apparent molecular weight of the  $\kappa$  light chains of DAV-1 (D<sub>L</sub>) was identical to that of the recombinant DT molecule (approx. 25 kDa). Western blot analyses showed that the DAV-1 mAb (D) and the DT molecules were recognized by an anti-mouse IgG antibody, while only the DT molecule was recognized by an anti-TNF- $\alpha$  polyclonal antibody.

DT molecules were capable of binding to immobilized penton base or Ad particles in an ELISA and elicited cytotoxicity against a TNF- $\alpha$  sensitive cell line,

-79-

MCF-7, indicating that the DT bifunctional molecule retains virus and cytokine receptor binding functions.

**Bifunctional molecules promote Ad-mediated gene delivery to  $\alpha v$  integrins**

- A first generation adenovirus vector containing a RSV-driven *LacZ* reporter gene with DT was preincubated at a ratio of 2 antibody molecules per RGD motif. This complex was then added to M21-L12 human melanoma cells, which do not express  $\alpha v$  integrins (Felding-Habermann *et al.* (1992) *J.Clin.Invest.* 89:2018-2022), but can support efficient virus binding (Wickham *et al.* (1993) *Cell* 73:309-319). Ad complexed with DT but not D alone, significantly increases Ad-mediated gene delivery to M21-L12 cells as measured by transgene expression at 48 hrs post-infection. Approximately 60% of cells incubated with Ad plus DT stained positive for  $\beta$ -galactosidase, compared to less than 3% of cells that had been incubated with virus alone or virus plus D. The increase in gene delivery by DT was not due to increased activation of the RSV LTR transgene promoter as a consequence of ligation of the TNF receptor, since M21-L12 cells that had been infected with adenovirus alone for three hours followed by addition of DT showed very little increase in gene delivery at 48 hours post-infection. This result indicates that bifunctional molecules increase adenovirus-mediated gene delivery by enhancing one or more steps associated with cell entry.

**DT molecules enhance Ad binding and internalization.**

- The following experiments showed that DT enhancement of gene delivery was associated with increased virus attachment to M21-L12 cells.  $^{125}\text{I}$ -labeled Ad particles alone or complexed with D or DT molecules were analyzed for binding to M21-L12 cells. Binding of  $^{125}\text{I}$ -Ad complexed with DT molecules was also examined in the absence or presence of an excess of recombinant Ad5 fiber protein, a polyclonal anti-human TNF- $\alpha$  antibody or a combination of the TNF- $\alpha$  antibody and fiber protein prior to addition to M21-L12 cells.  $^{125}\text{I}$ -labeled Ad was incubated with DT or D molecules prior to addition to M21-L12 cells and incubation on ice for 60 min. Unbound Ad was removed by washing and the cells were warmed to 37°C for varying times to allow virus internalization as measured by resistance to trypsinization. The results of these experiments

-80-

showed that pre-incubation of  $^{125}\text{I}$ -labeled Ad particles with DT but not with D molecules increased virus binding approximately 5-fold.

To investigate the molecules responsible for increased binding, competition experiments were performed. Ad-DT binding to cells was measured in the presence of a 50-fold excess of recombinant fiber protein or anti-TNF- $\alpha$  or a combination of these molecules. Either recombinant fiber or anti-TNF- $\alpha$  antibody alone was capable of blocking only 20-25% of Ad-DT binding to cells. In contrast, approximately 70% of binding could be inhibited by a combination of fiber and anti-TNF- $\alpha$ . These findings indicate that Ad-DT binding to cells is mediated by CAR-fiber internalization as well as TNF- $\alpha$ -receptor association. DT molecules potentiate internalization of  $^{125}\text{I}$ -labeled virus particles as measured by resistance to trypsin digestion.

As demonstrated in earlier studies (Wickham *et al.* (1993) *Cell* 73:309-319), because M21-L12 cells do not express  $\alpha\text{v}$  integrins, relatively low levels of adenovirus internalization by these cells occurs. DT molecules significantly increased the rate and extent of adenovirus internalization into these cells. These findings indicate that DT molecules enhance gene delivery by promoting virus binding as well as virus internalization.

DT enhancement of gene delivery is associated with PI3K activation.

Efficient Ad internalization via  $\alpha\text{v}$  integrins requires activation of PI3K, a key cellular signaling molecule. Experiments demonstrating that DT enhancement of gene delivery is mediated by PI3-kinase were performed. In these experiments, to show that DT enhancement of gene delivery also involves PI3K, M21-L12 melanoma cells were pretreated with the PI3K inhibitors wortmannin (30 nM) or LY292004 (20  $\mu\text{M}$ ) for 30 min prior to the addition of Ad.RSV. $\beta\text{gal}$  complexed with D or DT molecules. Ad mediated gene delivery was analyzed 48 hours post infection. The results show that wortmannin and LY294002 inhibited Ad-mediated gene delivery by approximately 70% and 50%, respectively, indicating that PI3K activity plays a major role in DT enhancement of gene delivery.

For further demonstration of the role of PI3K-dependent signaling in enhanced gene delivery, Ad-mediated gene delivery by other bifunctional

-81-

molecules whose cytokine/growth factor domains are known to activate PI3K was measured. Bifunctional molecules that target PI3K signaling pathways (DT = DAV-1-TNF; DE = DAV1-EGF; DI = DAV-1-IGF-1) also enhance Ad-mediated gene delivery to M21-L12 cells. DT, DE and DI molecules enhanced gene  
5 delivery by approximately 30, 10 and 5 fold respectively. Enhanced gene delivery by these molecules was also inhibited by pretreatment of cells with wortmannin. These findings further demonstrate that PI3K activation promotes Ad gene delivery.

#### EXAMPLE 6

##### 10 DEMONSTRATION OF TARGETED DELIVERY VIA GROWTH FACTOR RECEPTORS

Bifunctional molecules allow gene delivery by fiberless adenovirus particles

Fiberless adenovirus vector that cannot bind to CAR have been  
15 constructed (see Example 3 and also, e.g., copending U.S. application Serial No. 09/482,682 and U.S. application Serial No. 09/562,934; see, also Von Seggern *et al.* (1999) *J. Virol.* 73:1601-1608). The structure of these particles is nearly identical to that of wildtype virions. Fiberless particles alone showed almost no transgene delivery to SW480 epithelial cells, even though these cells express  
20 CAR and integrin  $\alpha\beta 5$  (Von Seggern *et al.* (1999) *J. Virol.* 73:1601-1608). To show that bifunctional molecules promote gene delivery by a fiberless Ad vector, CAR and  $\alpha\beta$  integrin-expressing SW480 cells were infected with a fiberless adenovirus or fiberless virus complexed with DT molecules at a ratio of 1 antibody molecule per viral particle. The reporter gene expression was examined  
25 by  $\beta$ -gal staining 48 hours post infection. The fiberless virus was incubated with bifunctional molecules before incubation with SW480 cells at 37° for 15 min. Ad-mediated gene delivery was examined 48 post infection.

The results demonstrated that DT molecules enhanced gene delivery of fiberless viruses. Fiberless particles complexed with DT or DI molecules  
30 exhibited increased gene delivery approximately 10-15, 3 and 5 fold, respectively, compared to the uncomplexed fiberless particles. These findings indicate that fiberless particles can be retargeted to cells via signal transducing antibodies.

-82-

Since modifications will be apparent to those of skill in this art, it is intended that this invention be limited only by the scope of the appended claims.

-83-

**CLAIMS:**

1. A bifunctional molecule, comprising:  
an antibody or antigen-binding portion thereof and a targeting  
agent, wherein:
  - 5 the antibody specifically binds to an antigen in a protein that binds to  $\alpha_v$  integrin; and
  - the targeting agent specifically binds to a cell surface protein that activates the phosphatidylinositol 3 (PI3K) signaling pathway.
2. The bifunctional molecule of claim 1, wherein the targeting agent  
10 or portion thereof triggers phosphatidylinositol-3-OH kinase (PI3K) activation.
3. The bifunctional molecule of claim 1, further comprising a linker that links the antibody or antigen-binding portion thereof to the targeting agent.
4. The bifunctional molecule of claim 1 that comprises a fusion  
protein.
- 15 5. The bifunctional molecule of claim 1 that comprises chemically conjugated polypeptides.
6. The bifunctional molecule of claim 2, wherein the linker is a single amino acid or a peptide.
7. The bifunctional molecule of claim 1, wherein the antibody  
20 comprises a heavy chain or a portion thereof sufficient for antigen-binding fused to the targeting agent.
8. The bifunctional molecule of claim 1, wherein the antibody portion is an Fab'2 fragment.
9. The bifunctional molecule of claim 1, wherein the antibody portion  
25 comprises a sufficient portion of the variable regions of the heavy and light chains for antigen recognition.
10. The bifunctional molecule of claim 1, wherein the antibody comprises the sequence of amino acids set forth in SEQ ID No. 2 or SEQ ID No. 6 or a sufficient portion thereof for antigen recognition.
- 30 11. The bifunctional molecule of claim 1, wherein the antibody comprises the sequence of amino acids set forth in SEQ ID No. 4 or a sufficient portion thereof for antigen recognition.



-84-

12. The bifunctional molecule of claim 1, wherein the antibody portion is an Fab fragment.

13. The bifunctional molecule of claim 10, wherein the nucleic acid encoding the antibody portion selected from

5 (a) the coding portion of the sequence of nucleotides set forth in SEQ ID No. 1 or SEQ ID No.5;

(b) a sequence of nucleotides that comprises one or more degenerate codons of (a); and

10 (c) a sequence of nucleotides that hybridizes along its full length under conditions of high stringency to a sufficient portion of (a) or (b) to encode an antigen-binding portion of the antibody.

14. The bifunctional molecule of claim 11, wherein the nucleic acid encoding the antibody portion is selected from

15 (a) the coding portion of the sequence of nucleotides set forth in SEQ ID No. 3;

(b) a sequence of nucleotides that comprises one or more degenerate codons of (a); and

20 (c) a sequence of nucleotides that hybridizes along its full length under conditions of high stringency to a sufficient portion of (a) or (b) to encode an antigen-binding portion of the antibody.

15. The bifunctional molecule of claim 1, comprising the sequence of amino acids set forth in any of SEQ ID Nos. 7-14 for specific binding to a targeted receptor.

25 16. The bifunctional molecule of claim 1, wherein the protein that binds to  $\alpha_v$  integrin is a viral protein or a bacterial protein that interacts with  $\alpha_v$  integrins for internalization of the respective virus or bacteria.

17. The bifunctional molecule of claim 16, wherein the virus is selected from adenovirus and adeno-associated virus.

30 18. The bifunctional molecule of claim 1, wherein the antibody or portion thereof specifically binds to the penton base of an adenovirus.

19. The bifunctional molecule of claim 1, wherein the antibody or portion thereof specifically binds to an antigen that includes an RGD motif.

-85-

20. The bifunctional molecule of claim 1, wherein the targeting agent comprises all or sufficient portion thereof of a protein that binds to G-protein coupled receptors, oncogene product receptors, hormone receptors or cytokine receptors that employ the PI3 signalling pathway for signal transduction,

5 wherein the sufficient portion thereof specifically binds to the cell surface receptor therefor and internalizes linked viral or bacterial particles.

21. The bifunctional molecule of claim 1, wherein the targeting agent comprises all or sufficient portion thereof of a protein that binds to G-protein coupled receptors that employ the PI3 signalling pathway for signal transduction,

10 wherein the sufficient portion thereof specifically binds to the cell surface receptor therefor and internalizes linked viral or bacterial particles.

22. The bifunctional molecule of claim 1, wherein the targeting agent comprises all or sufficient portion thereof of hormone or growth factor or cytokine,

15 wherein the sufficient portion thereof is specifically bind to the cell surface receptor therefor and internalizes linked viral or bacterial particles.

23. The bifunctional molecule of claim 6, wherein the targeting agent or portion thereof is a tumor necrosis factor (TNF), an fibroblast growth factor (FGF), an insulin-like growth factor (IGF) a colony stimulating factor (CSF),

20 insulin or a serum cell factor (SCF).

24. The bifunctional molecule of claim 6, wherein the targeting agent or portion thereof is insulin, IGF-1, TNF- $\alpha$ , SCF, CSF, a PDGF, an FGF, a heparin binding epidermal growth factor (HBEGF), a VEGF or dimer thereof.

25 25. The bifunctional molecule of claim 6, wherein the targeting agent or portion thereof is TNF- $\alpha$ , IGF-1, SCF or EGF.

26. The bifunctional molecule of claim 1, wherein targeted cell surface protein is selected from among a PDGF receptor, an IGF-1 receptor, an EGF receptor, a member of the FGF receptor family, a TNF receptor, a CSF-1 receptor, an insulin receptor, an IGF-1 receptor, an NGF receptor, an IL-2 receptor, an IL-3 receptor, an IL-4 receptor, an IgM receptor, a CD4 receptor, a CD2 receptor, a CD3/T cell receptor, a G protein linked thrombin receptor, an ATP receptor, and an fMLP receptor.

-86-

27. The bifunctional molecule of claim 1, wherein the targeted cell surface protein is selected from among tyrosine kinase receptors that, when activated, result in increased accumulation of PtdIns(3,4,5)P<sub>3</sub>, receptors associated with the src family non-receptor tyrosine kinases that stimulate PI3Ks  
5 phosphorylate phosphatidylinositol(3,4,5)P<sub>3</sub> (PtdIns(3,4,5)P<sub>3</sub>) accumulation.

28. An isolated nucleic acid molecule(s), comprising a sequence of nucleotides that encodes the bifunctional molecule of claim 1.

29. A targeted delivery vector, comprising:  
a bifunctional molecule of claim 1; and  
10 a viral or bacterial vector.

30. The targeted delivery vector of claim 29, wherein the gene delivery vector encodes a therapeutic product.

31. The targeted delivery vector of claim 29, wherein the vector is an adenovirus vector.

15 32. The targeted delivery vector of claim 29, wherein the vector is a fiberless adenovirus vector.

33. The targeted delivery vector of claim 29, wherein the bifunctional molecule and viral or bacterial vector are complexed by interaction of the antibody portion of the bifunctional molecule with a viral or bacterial surface  
20 protein.

34. The targeted delivery vector of claim 12, wherein the bifunctional molecule and viral or bacterial vector wherein the antibody portion of the bifunctional molecule is covalently linked to the viral or bacterial surface protein.

25 35. An isolated nucleic acid molecule, comprising a sequence of nucleotides encoding the bifunctional molecule of claim 1.

36. A combination, comprising:

a delivery vector for delivering gene products to targeted cells; and  
a bifunctional molecule of claim 1.

30 37. The combination of claim 36, wherein the bifunctional molecule and delivery vector for delivering gene products to targeted cells are complexed.

38. A method of targeted gene therapy, comprising administering a combination of claim 36.

-87-

39. A method of targeted gene therapy, comprising administering a combination of claim 37.

## SEQUENCE LISTING

&lt;110&gt; Nemerow, Glen R. et al.

&lt;120&gt; BIFUNCTIONAL MOLECULES AND VECTORS COMPLEXED THEREWITH FOR TARGETED GENE DELIVERY

&lt;130&gt; 22908-1228

&lt;140&gt; Unassigned

&lt;141&gt; 2000-07-10

&lt;160&gt; 33

&lt;170&gt; FastSEQ for Windows Version 4.0

&lt;210&gt; 1

&lt;211&gt; 1516

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (28)...(1395)

&lt;223&gt; DAV-1 heavy chain, penton base monoclonal antibody

&lt;400&gt; 1

cagacactga acacactgac tctaacc atg gga tgg agc tgg atc ttt ctc ttc 54  
 Met Gly Trp Ser Trp Ile Phe Leu Phe  
 1 5

ctc ctg tca gga act gca ggc gtc cac tct gag gtc cag ctt cag cag 102  
 Leu Leu Ser Gly Thr Ala Gly Val His Ser Glu Val Gln Leu Gln Gln 25  
 10 15 20

tca gga cct gag ctg gtg aaa cct ggg gcc tca gtg aag ata tcc tgc 150  
 Ser Gly Pro Glu Leu Val Lys Pro Gly Ala Ser Val Lys Ile Ser Cys 40  
 30 35

aag gct tct gga tac aca ttc act gac tac aac atg cac tgg gtg aag 198  
 Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr Asn Met His Trp Val Lys 55  
 45 50

cag agc cat gga aag agc ctt gag tgg att gga tat att tat cct tac 246  
 Gln Ser His Gly Lys Ser Leu Glu Trp Ile Gly Tyr Ile Tyr Pro Tyr 70  
 60 65

aaa ggt ggt act ggc tac aac cag aag ttc aag agc aag gcc aca ttg 294  
 Lys Gly Gly Thr Gly Tyr Asn Gln Lys Phe Lys Ser Lys Ala Thr Leu 85  
 75 80

aca aca gac agt tcc tcc aac aca gcc tac atg gag ctc cgc agc ctg 342  
 Thr Thr Asp Ser Ser Ser Asn Thr Ala Tyr Met Glu Leu Arg Ser Leu 105  
 90 95 100

aca tct gat gcc tct gca gtc tat tac tgt gca aga ggg att gct tac 390  
 Thr Ser Asp Ala Ser Ala Val Tyr Tyr Cys Ala Arg Gly Ile Ala Tyr 120  
 110 115

tgg ggc caa ggg act ctg gtc act gtc tct gca gcc aaa acg aca ccc 438  
 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala Ala Lys Thr Thr Pro 135  
 125 130

cca tct gtc tat cca ctg gcc cct gga tct gct gcc caa act aac tcc 486  
 Pro Ser Val Tyr Pro Leu Ala Pro Gly Ser Ala Ala Gln Thr Asn Ser 150  
 140 145

atg gtg acc ctg gga tgc ctg gtc aag ggc tat ttc cct gag cca gtg Met Val Thr Leu Gly Cys Leu Val Lys Gly Tyr Phe Pro Glu Pro Val 155 160 165	534
aca gtg acc tgg aac tct gga tcc ctg tcc agc ggt gtg cac acc ttc Thr Val Thr Trp Asn Ser Gly Ser Leu Ser Ser Gly Val His Thr Phe 170 175 180 185	582
cca gct gtc ctg cag tct gac ctc tac act ctg agc agc tca gtg act Pro Ala Val Leu Gln Ser Asp Leu Tyr Thr Leu Ser Ser Ser Val Thr 190 195 200	630
gtc ccc tcc agc acc tgg ccc agc gag acc gtc acc tgc aac gtt gcc Val Pro Ser Ser Thr Trp Pro Ser Glu Thr Val Thr Cys Asn Val Ala 205 210 215	678
cac ccg gcc agc agc acc aag gtg gac aag aaa att gtg ccc agg gat His Pro Ala Ser Ser Thr Lys Val Asp Lys Lys Ile Val Pro Arg Asp 220 225 230	726
tgt ggt tgt aag cct tgc ata tgt aca gtc cca gaa gta tca tct gtc Cys Gly Cys Lys Pro Cys Ile Cys Thr Val Pro Glu Val Ser Ser Val 235 240 245	765
ttc atc ttc ccc cca aag ccc aag gat gtg ctc acc att act ctg act Phe Ile Phe Pro Pro Lys Pro Lys Asp Val Leu Thr Ile Thr Leu Thr 250 255 260 265	822
cct aag gtc acg tgt gtt gtg gta gac atc agc aag gat gat ccc gag Pro Lys Val Thr Cys Val Val Val Asp Ile Ser Lys Asp Asp Pro Glu 270 275 280	870
gtc cag ttc agc tgg ttt gta gat gat gtg gag gtg cac aca gct cag Val Gln Phe Ser Trp Phe Val Asp Asp Val Glu Val His Thr Ala Gln 285 290 295	918
acg caa ccc cgg gag gag cag ttc aac agc act ttc cgc tca gtc agt Thr Gln Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg Ser Val Ser 300 305 310	966
gaa ctt ccc atc atg cac cag gac tgg ctc aat ggc aag gag ttc aaa Glu Leu Pro Ile Met His Gln Asp Trp Leu Asn Gly Lys Glu Phe Lys 315 320 325	1014
tgc agg gtc aac agt gca gct ttc cct gcc ccc atc gag aaa acc atc Cys Arg Val Asn Ser Ala Ala Phe Pro Ala Pro Ile Glu Lys Thr Ile 330 335 340 345	1062
tcc aaa acc aaa ggc aga cgg aag gct cca cag gtg tac acc att cca Ser Lys Thr Lys Gly Arg Pro Lys Ala Pro Gln Val Tyr Thr Ile Pro 350 355 360	1110
cct ccc aag gag cag atg gcc aag gat aaa gtc agt ctg acc tgc atg Pro Pro Lys Glu Gln Met Ala Lys Asp Lys Val Ser Leu Thr Cys Met 365 370 375	1158
ata aca gac ttc ttc cct gaa gac att act gtg gag tgg cag tgg aat Ile Thr Asp Phe Phe Pro Glu Asp Ile Thr Val Glu Trp Gln Trp Asn 380 385 390	1206
ggg cag cca gcg gag aac tac aag aac act cag ccc atc atg gac aca Gly Gln Pro Ala Glu Asn Tyr Lys Asn Thr Gln Pro Ile Met Asp Thr 395 400 405	1254
gat ggc tct tac ttc gtc tac agc aag ctc aat gtg cag aag agc aac Asp Gly Ser Tyr Phe Val Tyr Ser Lys Leu Asn Val Gln Lys Ser Asn 410 415 420 425	1302

tgg gag gca gga aat act ttc atc tgc tct gtg tta cat gag ggc ctg 1350  
 Trp Glu Ala Gly 430 Thr Phe Ile Cys Ser Val Leu His Glu Gly Leu 440

cac aac cac cat act gag aag agc ctc tcc cac tct cct ggt aaa 1395  
 His Asn His His Thr Glu Lys Ser Leu Ser His Ser Pro Gly Lys 445 455 455

tgateccagcgt gtccttgga g cccctctgggc ctacaggact ctgtcaccta cctccacccc 1455  
 tccctgtata aataaagcac cttagcactgc cttgggaccc tgcaataaaa aaaaaaaaaa 1515  
 a 1516

<210> 2  
 <211> 456  
 <212> PRT  
 <213> Mouse

<220>  
 <221> PEPTIDE  
 <222> (0)...(0)  
 <223> DAV-1 heavy chain, penton base monoclonal antibody

<400> 2  
 Met Gly Trp Ser Trp Ile Phe Leu Phe Leu Leu Ser Gly Thr Ala Gly  
 1 5 10 15  
 Val His Ser Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys  
 20 25 30  
 Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 35 40 45  
 Thr Asp Tyr Asn Met His Trp Val Lys Gln Ser His Gly Lys Ser Leu  
 50 55 60  
 Glu Trp Ile Gly Tyr Ile Tyr Pro Tyr Lys Gly Gly Thr Gly Tyr Asn  
 65 70 75 80  
 Gln Lys Phe Lys Ser Lys Ala Thr Leu Thr Thr Asp Ser Ser Ser Asn  
 85 90 95  
 Thr Ala Tyr Met Glu Leu Arg Ser Leu Thr Ser Asp Ala Ser Ala Val  
 100 105 110  
 Tyr Tyr Cys Ala Arg Gly Ile Ala Tyr Trp Gly Gln Gly Thr Leu Val  
 115 120 125  
 Thr Val Ser Ala Ala Lys Thr Thr Pro Pro Ser Val Tyr Pro Leu Ala  
 130 135 140  
 Pro Gly Ser Ala Ala Gln Thr Asn Ser Met Val Thr Leu Gly Cys Leu  
 145 150 155 160  
 Val Lys Gly Tyr Phe Pro Glu Pro Val Thr Val Thr Trp Asn Ser Gly  
 165 170 175  
 Ser Leu Ser Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Asp  
 180 185 190  
 Leu Tyr Thr Leu Ser Ser Ser Val Thr Val Pro Ser Ser Thr Trp Pro  
 195 200 205  
 Ser Glu Thr Val Thr Cys Asn Val Ala His Pro Ala Ser Ser Thr Lys  
 210 215 220  
 Val Asp Lys Lys Ile Val Pro Arg Asp Cys Gly Cys Lys Pro Cys Ile  
 225 230 235 240  
 Cys Thr Val Pro Glu Val Ser Ser Val Phe Ile Phe Pro Pro Lys Pro  
 245 250 255  
 Lys Asp Val Leu Thr Ile Thr Leu Thr Pro Lys Val Thr Cys Val Val  
 260 265 270  
 Val Asp Ile Ser Lys Asp Asp Pro Glu Val Gln Phe Ser Trp Phe Val  
 275 280 285  
 Asp Asp Val Glu Val His Thr Ala Gln Thr Gln Pro Arg Glu Glu Gln  
 290 295 300  
 Phe Asn Ser Thr Phe Arg Ser Val Ser Glu Leu Pro Ile Met His Gln  
 305 310 315 320  
 Asp Trp Leu Asn Gly Lys Glu Phe Lys Cys Arg Val Asn Ser Ala Ala  
 325 330 335  
 Phe Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Arg Pro

```

          340          345          350
Lys Ala Pro Gln Val Tyr Thr Ile Pro Pro Pro Lys Glu Gln Met Ala
          355          360          365
Lys Asp Lys Val Ser Leu Thr Cys Met Ile Thr Asp Phe Phe Pro Glu
          370          375          380
Asp Ile Thr Val Glu Trp Gln Trp Asn Gly Gln Pro Ala Glu Asn Tyr
          385          390          395          400
Lys Asn Thr Gln Pro Ile Met Asp Thr Asp Gly Ser Tyr Phe Val Tyr
          405          410          415
Ser Lys Leu Asn Val Gln Lys Ser Asn Trp Glu Ala Gly Asn Thr Phe
          420          425          430
Ile Cys Ser Val Leu His Glu Gly Leu His Asn His His Thr Glu Lys
          435          440          445
Ser Leu Ser His Ser Pro Gly Lys
          450          455

```

&lt;210&gt; 3

&lt;211&gt; 831

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (13)...(726)

&lt;223&gt; DAV-1 light chain, penton base monoclonal antibody

&lt;400&gt; 3

```

aagcttaccg cc atg gag aca gac aca atc ctg cta tgg gtg ctg ctg ctc      51
      Met Glu Thr Asp Thr Ile Leu Leu Trp Val Leu Leu Leu
          1                      5                      10

tgg gtt cca ggc tcc act ggt gac att gtg ctg acc caa tct cca gct      99
Trp Val Pro Gly Ser Thr Gly Asp Ile Val Leu Thr Gln Ser Pro Ala
          15                      20                      25

tct ttg gct gtg tct cta ggg cag agg gcc acc atc tcc tgc aag gcc      147
Ser Leu Ala Val Ser Leu Gly Gln Arg Ala Thr Ile Ser Cys Lys Ala
          30                      35                      40                      45

agc caa agt gtt gat tat gat ggt gat agt tat atg aac tgg tac caa      195
Ser Gln Ser Val Asp Tyr Asp Gly Asp Ser Tyr Met Asn Trp Tyr Gln
          50                      55                      60

cag aaa cca gga cag cca ccc aaa ctc ctc atc tat gct gca tcc aat      243
Gln Lys Pro Gly Gln Pro Pro Lys Leu Ile Tyr Ala Ala Ser Asn
          65                      70                      75

tta gaa tct ggg atc cca gcc agg ttt agt ggc agt ggg tct ggg aca      291
Leu Glu Ser Gly Ile Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr
          80                      85                      90

gac ttc acc ctc aac atc cat cct gtg gag gag gag gat gct gca acc      339
Asp Phe Thr Leu Asn Ile His Pro Val Glu Glu Glu Asp Ala Ala Thr
          95                      100                      105

tat tac tgt cag caa act aat gag gat ccg tgg acg ttc ggt gga ggc      387
Tyr Tyr Cys Gln Gln Thr Asn Glu Asp Pro Trp Thr Phe Gly Gly Gly
          110                      115                      120                      125

acc aag ctg gaa atc aaa cgg gct gat gct gca cca act gta tcc atc      435
Thr Lys Leu Glu Ile Lys Arg Ala Asp Ala Ala Pro Thr Val Ser Ile
          130                      135                      140

ttc cca cca tcc agt gag cag tta aca tct gga ggt gcc tca gtc gtg      483
Phe Pro Pro Ser Ser Glu Gln Leu Thr Ser Gly Gly Ala Ser Val Val
          145                      150                      155

```



tgc ttc ttg aac aac ttc tac ccc aaa gac atc aat gtc aag tgg aag 531  
 Cys Phe Leu Asn Asn Phe Tyr Pro Lys Asp Ile Asn Val Lys Trp Lys  
 160 165 170

att gat ggc agt gaa cga caa aat ggc gtc ctg aac agt tgg act gat 579  
 Ile Asp Gly Ser Glu Arg Gln Asn Gly Val Leu Asn Ser Trp Thr Asp  
 175 180 185

cag gac agc aaa gac agc acc tac agc atg agc agc acc ctc acg ttg 627  
 Gln Asp Ser Lys Asp Ser Thr Tyr Ser Met Ser Ser Thr Leu Thr Leu  
 190 195 200 205

acc aag gac gag tat gaa cga cat aac agc tat acc tgt gag gcc act 675  
 Thr Lys Asp Glu Tyr Glu Arg His Asn Ser Tyr Thr Cys Glu Ala Thr  
 210 215 220

cac aag aca tca act tca ccc att gtc aag agc ttc aac agg aat gag 723  
 His Lys Thr Ser Thr Ser Pro Ile Val Lys Ser Phe Asn Arg Asn Glu  
 225 230 235

tgt tagagacaaa ggtcctgaga cgccaccacc agctccccag ctccatccta 776  
 Cys

tcttcccttc taaggctcttg gaggcttcct cgagcggtaa agggcgaatt ccage 831

<210> 4  
 <211> 238  
 <212> PRT  
 <213> Mouse

<220>  
 <221> PEPTIDE  
 <222> (0)...(0)  
 <223> DAV-1 light chain, penton base monoclonal antibody

<400> 4  
 Met Glu Thr Asp Thr Ile Leu Leu Trp Val Leu Leu Leu Trp Val Pro  
 1 5 10 15  
 Gly Ser Thr Gly Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala  
 20 25 30  
 Val Ser Leu Gly Gln Arg Ala Thr Ile Ser Cys Lys Ala Ser Gln Ser  
 35 40 45  
 Val Asp Tyr Asp Gly Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro  
 50 55 60  
 Gly Gln Pro Pro Lys Leu Ile Tyr Ala Ala Ser Asn Leu Glu Ser  
 65 70 75 80  
 Gly Ile Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr  
 85 90 95  
 Leu Asn Ile His Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys  
 100 105 110  
 Gln Gln Thr Asn Glu Asp Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu  
 115 120 125  
 Glu Ile Lys Arg Ala Asp Ala Ala Pro Thr Val Ser Ile Phe Pro Pro  
 130 135 140  
 Ser Ser Glu Gln Leu Thr Ser Gly Gly Ala Ser Val Val Cys Phe Leu  
 145 150 155 160  
 Asn Asn Phe Tyr Pro Lys Asp Ile Asn Val Lys Trp Lys Ile Asp Gly  
 165 170 175  
 Ser Glu Arg Gln Asn Gly Val Leu Asn Ser Trp Thr Asp Gln Asp Ser  
 180 185 190  
 Lys Asp Ser Thr Tyr Ser Met Ser Ser Thr Leu Thr Leu Thr Lys Asp  
 195 200 205  
 Glu Tyr Glu Arg His Asn Ser Tyr Thr Cys Glu Ala Thr His Lys Thr  
 210 215 220  
 Ser Thr Ser Pro Ile Val Lys Ser Phe Asn Arg Asn Glu Cys  
 225 230 235

<210> 5  
 <211> 1314  
 <212> DNA  
 <213> Mouse

<220>  
 <221> CDS  
 <222> (0)...(1314)  
 <223> Portion of DAV-1 heavy chain used for fusion protein  
 bifunctional antibody

```

<400> 5
atg gga tgg agc tgg atc ttt ctc ttc ctc ctg tca gga act gca ggc      48
Met Gly Trp Ser Trp Ile Phe Leu Phe Leu Leu Ser Gly Thr Ala Gly
  1                               5                               10                               15

gtc cac tct gag gtc cag ctt cag cag tca gga cct gag ctg gtg aaa      96
Val His Ser Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys
                20                               25                               30

cct ggg gcc tca gtg aag ata tcc tgc aag gct tct gga tac aca ttc      144
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe
                35                               40                               45

act gac tac aac atg cac tgg gtg aag cag agc cat gga aag agc ctt      192
Thr Asp Tyr Asn Met His Trp Val Lys Gln Ser His Gly Lys Ser Leu
                50                               55                               60

gag tgg att gga tat att tat cct tac aaa ggt ggt act ggc tac aac      240
Glu Trp Ile Gly Tyr Ile Tyr Pro Tyr Lys Gly Gly Thr Gly Tyr Asn
  65                               70                               75                               80

cag aag ttc aag agc aag gcc aca ttg aca aca gac agt tcc tcc aac      288
Gln Lys Phe Lys Ser Lys Ala Thr Leu Thr Thr Asp Ser Ser Ser Asn
                85                               90                               95

aca gcc tac atg gag ctc cgc agc ctg aca tct gat gcc tct gca gtc      336
Thr Ala Tyr Met Glu Leu Arg Ser Leu Thr Ser Asp Ala Ser Ala Val
                100                              105                              110

tat tac tgt gca aga ggg att gct tac tgg ggc caa ggg act ctg gtc      384
Tyr Tyr Cys Ala Arg Gly Ile Ala Tyr Trp Gly Gln Gly Thr Leu Val
                115                              120                              125

act gtc tct gca gcc aaa acg aca ccc cca tct gtc tat cca ctg gcc      432
Thr Val Ser Ala Ala Lys Thr Thr Pro Pro Ser Val Tyr Pro Leu Ala
                130                              135                              140

cct gga tct gct gcc caa act aac tcc atg gtg acc ctg gga tgc ctg      480
Pro Gly Ser Ala Ala Gln Thr Asn Ser Met Val Thr Leu Gly Cys Leu
                145                              150                              155                              160

gtc aag ggc tat ttc cct gag cca gtg aca gtg acc tgg aac tct gga      528
Val Lys Gly Tyr Phe Pro Glu Pro Val Thr Val Thr Trp Asn Ser Gly
                165                              170                              175

tcc ctg tcc agc ggt gtg cac acc ttc cca gct gtc ctg cag tct gac      576
Ser Leu Ser Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Asp
                180                              185                              190

ctc tac act ctg agc agc tca gtg act gtc ccc tcc agc acc tgg ccc      624
Leu Tyr Thr Leu Ser Ser Ser Val Thr Val Pro Ser Ser Thr Trp Pro
                195                              200                              205

agc gag acc gtc acc tgc aac gtt gcc cac ccg gcc agc agc acc aag      672
Ser Glu Thr Val Thr Cys Asn Val Ala His Pro Ala Ser Ser Thr Lys

```

210	215	220	
gtg gac aag aaa att Val Asp Lys Lys Ile 225	gtg ccc agg gat tgt Val Pro Arg Asp Cys 230	ggt tgt aag cct tgc ata Gly Cys Lys Pro Cys Ile 235 240	720
tgt aca gtc cca gaa gta tca tct gtc ttc atc ttc ccc cca aag ccc Cys Thr Val Pro Glu Val Ser Ser Val Phe Ile Phe Pro Pro Lys Pro 245 250 255			768
aag gat gtg ctc acc att act ctg act cct aag gtc acg tgt gtt gtg Lys Asp Val Leu Thr Ile Thr Leu Thr Pro Lys Val Thr Cys Val Val 260 265 270			816
gta gac atc agc aag gat gat ccc gag gtc cag ttc agc tgg ttt gta Val Asp Ile Ser Lys Asp Asp Pro Glu Val Gln Phe Ser Trp Phe Val 275 280 285			864
gat gat gtg gag gtg cac aca gct cag acg caa ccc cgg gag gag cag Asp Asp Val Glu Val His Thr Ala Gln Thr Gln Pro Arg Glu Glu Gln 290 295 300			912
ttc aac agc act ttc cgc tca gtc agt gaa ctt ccc atc atg cac cag Phe Asn Ser Thr Phe Arg Ser Val Ser Glu Leu Pro Ile Met His Gln 305 310 315 320			960
gac tgg ctc aat ggc aag gag ttc aaa tgc agg gtc aac agt gca gct Asp Trp Leu Asn Gly Lys Glu Phe Lys Cys Arg Val Asn Ser Ala Ala 325 330 335			1008
ttc cct gcc ccc atc gag aaa acc atc tcc aaa acc aaa ggc aga ccg Phe Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Arg Pro 340 345 350			1056
aag gct cca cag gtg tac acc att cca cct ccc aag gag cag atg gcc Lys Ala Pro Gln Val Tyr Thr Ile Pro Pro Pro Lys Glu Gln Met Ala 355 360 365			1104
aag gat aaa gtc agt ctg acc tgc atg ata aca gac ttc ttc cct gaa Lys Asp Lys Val Ser Leu Thr Cys Met Ile Thr Asp Phe Phe Pro Glu 370 375 380			1152
gac att act gtg gag tgg cag tgg aat ggg cag cca gcg gag aac tac Asp Ile Thr Val Glu Trp Gln Trp Asn Gly Gln Pro Ala Glu Asn Tyr 385 390 395 400			1200
aag aac act cag ccc atc atg gac aca gat ggc tct tac ttc gtc tac Lys Asn Thr Gln Pro Ile Met Asp Thr Asp Gly Ser Tyr Phe Val Tyr 405 410 415			1248
agc aag ctc aat gtg cag aag agc aac tgg gag gca gga aat act ttc Ser Lys Leu Asn Val Gln Lys Ser Asn Trp Glu Ala Gly Asn Thr Phe 420 425 430			1296
atc tgc tct gtg tta cat Ile Cys Ser Val Leu His 435			1314

<210> 6  
 <211> 438  
 <212> PRT  
 <213> Mouse

<220>  
 <221> PEPTIDE  
 <222> (0)....(0)

<223> Portion of DAV-1 heavy chain used for fusion protein  
bifunctional antibody

<400> 6

```

Met Gly Trp Ser Trp Ile Phe Leu Phe Leu Leu Ser Gly Thr Ala Gly
 1      5      10      15
Val His Ser Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys
 20      25      30
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe
 35      40      45
Thr Asp Tyr Asn Met His Trp Val Lys Gln Ser His Gly Lys Ser Leu
 50      55      60
Glu Trp Ile Gly Tyr Ile Tyr Pro Tyr Lys Gly Gly Thr Gly Tyr Asn
 65      70      75
Gln Lys Phe Lys Ser Lys Ala Thr Leu Thr Thr Asp Ser Ser Ser Asn
 85      90      95
Thr Ala Tyr Met Glu Leu Arg Ser Leu Thr Ser Asp Ala Ser Ala Val
100      105      110
Tyr Tyr Cys Ala Arg Gly Ile Ala Tyr Trp Gly Gln Gly Thr Leu Val
115      120      125
Thr Val Ser Ala Ala Lys Thr Thr Pro Pro Ser Val Tyr Pro Leu Ala
130      135      140
Pro Gly Ser Ala Ala Gln Thr Asn Ser Met Val Thr Leu Gly Cys Leu
145      150      155
Val Lys Gly Tyr Phe Pro Glu Pro Val Thr Val Thr Trp Asn Ser Gly
165      170      175
Ser Leu Ser Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Asp
180      185      190
Leu Tyr Thr Leu Ser Ser Ser Val Thr Val Pro Ser Ser Thr Trp Pro
195      200      205
Ser Glu Thr Val Thr Cys Asn Val Ala His Pro Ala Ser Ser Thr Lys
210      215      220
Val Asp Lys Lys Ile Val Pro Arg Asp Cys Gly Cys Lys Pro Cys Ile
225      230      235
Cys Thr Val Pro Glu Val Ser Ser Val Phe Ile Phe Pro Pro Lys Pro
245      250      255
Lys Asp Val Leu Thr Ile Thr Leu Thr Pro Lys Val Thr Cys Val Val
260      265      270
Val Asp Ile Ser Lys Asp Asp Pro Glu Val Gln Phe Ser Trp Phe Val
275      280      285
Asp Asp Val Glu Val His Thr Ala Gln Thr Gln Pro Arg Glu Glu Gln
290      295      300
Phe Asn Ser Thr Phe Arg Ser Val Ser Glu Leu Pro Ile Met His Gln
305      310      315
Asp Trp Leu Asn Gly Lys Glu Phe Lys Cys Arg Val Asn Ser Ala Ala
325      330      335
Phe Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Arg Pro
340      345      350
Lys Ala Pro Gln Val Tyr Thr Ile Pro Pro Pro Lys Glu Gln Met Ala
355      360      365
Lys Asp Lys Val Ser Leu Thr Cys Met Ile Thr Asp Phe Phe Pro Glu
370      375      380
Asp Ile Thr Val Glu Trp Gln Trp Asn Gly Gln Pro Ala Glu Asn Tyr
385      390      395
Lys Asn Thr Gln Pro Ile Met Asp Thr Asp Gly Ser Tyr Phe Val Tyr
405      410      415
Ser Lys Leu Asn Val Gln Lys Ser Asn Trp Glu Ala Gly Asn Thr Phe
420      425      430
Ile Cys Ser Val Leu His
435

```

<210> 7

<211> 157

<212> PRT

<213> Human

<220>  
 <221> PEPTIDE  
 <222> (0)...(0)  
 <223> Tumor necrosis factor-alpha (TNF alpha, mature peptide)

<400> 7  
 Val Arg Ser Ser Ser Arg Thr Pro Ser Asp Lys Pro Val Ala His Val  
 1 5 10 15  
 Val Ala Asn Pro Gln Ala Glu Gly Gln Leu Gln Trp Leu Asn Arg Arg  
 20 25 30  
 Ala Asn Ala Leu Leu Ala Asn Gly Val Glu Leu Arg Asp Asn Gln Leu  
 35 40 45  
 Val Val Pro Ser Glu Gly Leu Tyr Leu Ile Tyr Ser Gln Val Leu Phe  
 50 55 60  
 Lys Gly Gln Gly Cys Pro Ser Thr His Val Leu Leu Thr His Thr Ile  
 65 70 75 80  
 Ser Arg Ile Ala Val Ser Tyr Gln Thr Lys Val Asn Leu Leu Ser Ala  
 85 90 95  
 Ile Lys Ser Pro Cys Gln Arg Glu Thr Pro Glu Gly Ala Glu Ala Lys  
 100 105 110  
 Pro Trp Tyr Glu Pro Ile Tyr Leu Gly Gly Val Phe Gln Leu Glu Lys  
 115 120 125  
 Gly Asp Arg Leu Ser Ala Glu Ile Asn Arg Pro Asp Tyr Leu Asp Phe  
 130 135 140  
 Ala Glu Ser Gly Gln Val Tyr Phe Gly Ile Ile Ala Leu  
 145 150 155

<210> 8  
 <211> 70  
 <212> PRT  
 <213> Human

<220>  
 <221> PEPTIDE  
 <222> (0)...(0)  
 <223> Human Insulin-like Growth Factor 1 sequence  
 (IGF-1, mature peptide)

<400> 8  
 Gly Pro Glu Thr Leu Cys Gly Ala Glu Leu Val Asp Ala Leu Gln Phe  
 1 5 10 15  
 Val Cys Gly Asp Arg Gly Phe Tyr Phe Asn Lys Pro Thr Gly Tyr Gly  
 20 25 30  
 Ser Ser Ser Arg Arg Ala Pro Gln Thr Gly Ile Val Asp Glu Cys Cys  
 35 40 45  
 Phe Arg Ser Cys Asp Leu Arg Arg Leu Glu Met Tyr Cys Ala Pro Leu  
 50 55 60  
 Lys Pro Ala Lys Ser Ala  
 65 70

<210> 9  
 <211> 53  
 <212> PRT  
 <213> Human

<220>  
 <221> PEPTIDE  
 <222> (0)...(0)  
 <223> Epidermal Growth Factor (EGF, mature peptide)

<400> 9  
 Asn Ser Asp Ser Glu Cys Pro Leu Ser His Asp Gly Tyr Cys Leu His  
 1 5 10 15  
 Asp Gly Val Cys Met Tyr Ile Glu Ala Leu Asp Lys Tyr Ala Cys Asn  
 20 25 30  
 Cys Val Val Gly Tyr Ile Gly Glu Arg Cys Gln Tyr Arg Asp Leu Lys

35  
Trp Trp Glu Leu Arg  
50

40

45

<210> 10  
<211> 164  
<212> PRT  
<213> Human

<220>  
<221> PEPTIDE  
<222> (0)...(0)  
<223> Stem Cell Factor (SCF, mature peptide)

<400> 10  
Glu Gly Ile Cys Arg Asn Arg Val Thr Asn Asn Val Lys Asp Val Thr  
1 5 10 15  
Lys Leu Val Ala Asn Leu Pro Lys Asp Tyr Met Ile Thr Leu Lys Tyr  
20 25 30  
Val Pro Gly Met Asp Val Leu Pro Ser His Cys Trp Ile Ser Glu Met  
35 40 45  
Val Val Gln Leu Ser Asp Ser Leu Thr Asp Leu Leu Asp Lys Phe Ser  
50 55 60  
Asn Ile Ser Glu Gly Leu Ser Asn Tyr Ser Ile Ile Asp Lys Leu Val  
65 70 75 80  
Asn Ile Val Asp Asp Leu Val Glu Cys Val Lys Glu Asn Ser Ser Lys  
85 90 95  
Asp Leu Lys Lys Ser Phe Lys Ser Pro Glu Pro Arg Leu Phe Thr Pro  
100 105 110  
Glu Glu Phe Phe Arg Ile Phe Asn Arg Ser Ile Asp Ala Phe Lys Asp  
115 120 125  
Phe Val Val Ala Ser Glu Thr Ser Asp Cys Val Val Ser Ser Thr Leu  
130 135 140  
Ser Pro Glu Lys Asp Ser Arg Val Ser Val Thr Lys Pro Phe Met Leu  
145 150 155 160  
Pro Pro Val Ala

<210> 11  
<211> 597  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Fusion protein with N-terminal portion of DAV-1 heavy chain  
and TNF alpha mature peptide

<400> 11  
Met Gly Trp Ser Trp Ile Phe Leu Phe Leu Leu Ser Gly Thr Ala Gly  
1 5 10 15  
Val His Ser Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys  
20 25 30  
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
35 40 45  
Thr Asp Tyr Asn Met His Trp Val Lys Gln Ser His Gly Lys Ser Leu  
50 55 60  
Glu Trp Ile Gly Tyr Ile Tyr Pro Tyr Lys Gly Gly Thr Gly Tyr Asn  
65 70 75 80  
Gln Lys Phe Lys Ser Lys Ala Thr Leu Thr Asp Ser Ser Ser Asn  
85 90 95  
Thr Ala Tyr Met Glu Leu Arg Ser Leu Thr Ser Asp Ala Ser Ala Val  
100 105 110  
Tyr Tyr Cys Ala Arg Gly Ile Ala Tyr Trp Gly Gln Gly Thr Leu Val  
115 120 125  
Thr Val Ser Ala Ala Lys Thr Thr Pro Pro Ser Val Tyr Pro Leu Ala  
130 135 140  
Pro Gly Ser Ala Ala Gln Thr Asn Ser Met Val Thr Leu Gly Cys Leu

145	Val	Lys	Gly	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Thr	Trp	Asn	Ser	Gly	160
					165	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Asp	175
	Ser	Leu	Ser	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Asp	190
					180	Ser	Ser	Ser	Val	Thr	Val	Pro	Ser	Ser	Thr	Trp	Pro
	Leu	Tyr	Thr	Leu	Ser	Ser	Ser	Val	Thr	Val	Pro	Ser	Ser	Ser	Thr	Trp	Pro
					195	Thr	Cys	Asn	Val	Ala	His	Pro	Ala	Ser	Ser	Thr	Lys
	Ser	Glu	Thr	Val	Thr	Cys	Asn	Val	Ala	His	Pro	Ala	Ser	Ser	Ser	Thr	Lys
					210	Val	Pro	Arg	Asp	Cys	Gly	Cys	Lys	Pro	Cys	Ile	
	Val	Asp	Lys	Lys	Ile	Val	Pro	Arg	Asp	Cys	Gly	Cys	Lys	Pro	Cys	Ile	
					225	Cys	Thr	Val	Pro	Glu	Val	Ser	Ser	Val	Phe	Ile	Phe
	Cys	Thr	Val	Pro	Glu	Val	Ser	Ser	Val	Phe	Ile	Phe	Pro	Pro	Lys	Pro	
					245	Lys	Asp	Val	Leu	Thr	Ile	Thr	Leu	Thr	Pro	Lys	Val
	Lys	Asp	Val	Leu	Thr	Ile	Thr	Leu	Thr	Pro	Lys	Val	Thr	Cys	Val	Val	
					260	Val	Asp	Ile	Ser	Lys	Asp	Asp	Pro	Glu	Val	Gln	Phe
	Val	Asp	Ile	Ser	Lys	Asp	Asp	Pro	Glu	Val	Gln	Phe	Ser	Trp	Phe	Val	
					275	Asp	Asp	Val	Glu	Val	His	Thr	Ala	Gln	Thr	Gln	Pro
	Asp	Asp	Val	Glu	Val	His	Thr	Ala	Gln	Thr	Gln	Pro	Arg	Glu	Glu	Gln	
					290	Phe	Asn	Ser	Thr	Phe	Arg	Ser	Val	Ser	Glu	Leu	Pro
	Phe	Asn	Ser	Thr	Phe	Arg	Ser	Val	Ser	Glu	Leu	Pro	Ile	Met	His	Gln	
					305	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Phe	Lys	Cys	Arg	Val
	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Phe	Lys	Cys	Arg	Val	Asn	Ser	Ala	Ala	
					325	Phe	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Thr
	Phe	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Thr	Lys	Gly	Arg	Pro	
					340	Lys	Ala	Pro	Gln	Val	Tyr	Thr	Ile	Pro	Pro	Pro	Lys
	Lys	Ala	Pro	Gln	Val	Tyr	Thr	Ile	Pro	Pro	Pro	Lys	Glu	Gln	Met	Ala	
					355	Lys	Asp	Lys	Val	Ser	Leu	Thr	Cys	Met	Ile	Thr	Asp
	Lys	Asp	Lys	Val	Ser	Leu	Thr	Cys	Met	Ile	Thr	Asp	Phe	Phe	Pro	Glu	
					370	Asp	Ile	Thr	Val	Glu	Trp	Gln	Trp	Asn	Gly	Gln	Pro
	Asp	Ile	Thr	Val	Glu	Trp	Gln	Trp	Asn	Gly	Gln	Pro	Ala	Glu	Asn	Tyr	
					385	Lys	Asn	Thr	Gln	Pro	Ile	Met	Asp	Thr	Asp	Gly	Ser
	Lys	Asn	Thr	Gln	Pro	Ile	Met	Asp	Thr	Asp	Gly	Ser	Tyr	Phe	Val	Tyr	
					405	Ser	Lys	Leu	Asn	Val	Gln	Lys	Ser	Asn	Trp	Glu	Ala
	Ser	Lys	Leu	Asn	Val	Gln	Lys	Ser	Asn	Trp	Glu	Ala	Gly	Asn	Thr	Phe	
					420	Ile	Cys	Ser	Val	Leu	His	Glu	Phe	Val	Arg	Ser	Ser
	Ile	Cys	Ser	Val	Leu	His	Glu	Phe	Val	Arg	Ser	Ser	Ser	Arg	Thr	Pro	
					435	Ser	Asp	Lys	Pro	Val	Ala	His	Val	Val	Ala	Asn	Pro
	Ser	Asp	Lys	Pro	Val	Ala	His	Val	Val	Ala	Asn	Pro	Gln	Ala	Glu	Gly	
					450	Gln	Leu	Gln	Trp	Leu	Asn	Arg	Arg	Ala	Asn	Ala	Leu
	Gln	Leu	Gln	Trp	Leu	Asn	Arg	Arg	Ala	Asn	Ala	Leu	Leu	Ala	Asn	Gly	
					465	Val	Glu	Leu	Arg	Asp	Asn	Gln	Leu	Val	Val	Pro	Ser
	Val	Glu	Leu	Arg	Asp	Asn	Gln	Leu	Val	Val	Pro	Ser	Glu	Gly	Leu	Tyr	
					485	Leu	Ile	Tyr	Ser	Gln	Val	Leu	Phe	Lys	Gly	Gln	Gly
	Leu	Ile	Tyr	Ser	Gln	Val	Leu	Phe	Lys	Gly	Gln	Gly	Cys	Pro	Ser	Thr	
					500	His	Val	Leu	Leu	Thr	His	Thr	Ile	Ser	Arg	Ile	Ala
	His	Val	Leu	Leu	Thr	His	Thr	Ile	Ser	Arg	Ile	Ala	Val	Ser	Tyr	Gln	
					515	Thr	Lys	Val	Asn	Leu	Leu	Ser	Ala	Ile	Lys	Ser	Pro
	Thr	Lys	Val	Asn	Leu	Leu	Ser	Ala	Ile	Lys	Ser	Pro	Cys	Gln	Arg	Glu	
					530	Thr	Pro	Glu	Gly	Ala	Glu	Ala	Lys	Pro	Trp	Tyr	Glu
	Thr	Pro	Glu	Gly	Ala	Glu	Ala	Lys	Pro	Trp	Tyr	Glu	Pro	Ile	Tyr	Leu	
					545	Gly	Gly	Val	Phe	Gln	Leu	Glu	Lys	Gly	Asp	Arg	Leu
	Gly	Gly	Val	Phe	Gln	Leu	Glu	Lys	Gly	Asp	Arg	Leu	Ser	Ala	Glu	Ile	
					565	Asn	Arg	Pro	Asp	Tyr	Leu	Asp	Phe	Ala	Glu	Ser	Gly
	Asn	Arg	Pro	Asp	Tyr	Leu	Asp	Phe	Ala	Glu	Ser	Gly	Gln	Val	Tyr	Phe	
					580	Gly	Ile	Ile	Ala	Leu							
	Gly	Ile	Ile	Ala	Leu												
					595												

&lt;210&gt; 12

&lt;211&gt; 510

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Fusion protein with N-terminal portion of DAV-1 heavy chain and IGF-1 mature peptide

<400> 12  
 Met Gly Trp Ser Trp Ile Phe Leu Phe Leu Leu Ser Gly Thr Ala Gly  
 1 5 10 15  
 Val His Ser Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys  
 20 25 30  
 Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 35 40 45  
 Thr Asp Tyr Asn Met His Trp Val Lys Gln Ser His Gly Lys Ser Leu  
 50 55 60  
 Glu Trp Ile Gly Tyr Ile Tyr Pro Tyr Lys Gly Gly Thr Gly Tyr Asn  
 65 70 75 80  
 Gln Lys Phe Lys Ser Lys Ala Thr Leu Thr Thr Asp Ser Ser Ser Asn  
 85 90 95  
 Thr Ala Tyr Met Glu Leu Arg Ser Leu Thr Ser Asp Ala Ser Ala Val  
 100 105 110  
 Tyr Tyr Cys Ala Arg Gly Ile Ala Tyr Trp Gly Gln Gly Thr Leu Val  
 115 120 125  
 Thr Val Ser Ala Ala Lys Thr Thr Pro Pro Ser Val Tyr Pro Leu Ala  
 130 135 140  
 Pro Gly Ser Ala Ala Gln Thr Asn Ser Met Val Thr Leu Gly Cys Leu  
 145 150 155 160  
 Val Lys Gly Tyr Phe Pro Glu Pro Val Thr Val Thr Trp Asn Ser Gly  
 165 170 175  
 Ser Leu Ser Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Asp  
 180 185 190  
 Leu Tyr Thr Leu Ser Ser Ser Val Thr Val Pro Ser Ser Thr Trp Pro  
 195 200 205  
 Ser Glu Thr Val Thr Cys Asn Val Ala His Pro Ala Ser Ser Thr Lys  
 210 215 220  
 Val Asp Lys Lys Ile Val Pro Arg Asp Cys Gly Cys Lys Pro Cys Ile  
 225 230 235 240  
 Cys Thr Val Pro Glu Val Ser Ser Val Phe Ile Phe Pro Pro Lys Pro  
 245 250 255  
 Lys Asp Val Leu Thr Ile Thr Leu Thr Pro Lys Val Thr Cys Val Val  
 260 265 270  
 Val Asp Ile Ser Lys Asp Asp Pro Glu Val Gln Phe Ser Trp Phe Val  
 275 280 285  
 Asp Asp Val Glu Val His Thr Ala Gln Thr Gln Pro Arg Glu Glu Gln  
 290 295 300  
 Phe Asn Ser Thr Phe Arg Ser Val Ser Glu Leu Pro Ile Met His Gln  
 305 310 315 320  
 Asp Trp Leu Asn Gly Lys Glu Phe Lys Cys Arg Val Asn Ser Ala Ala  
 325 330 335  
 Phe Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Arg Pro  
 340 345 350  
 Lys Ala Pro Gln Val Tyr Thr Ile Pro Pro Pro Lys Glu Gln Met Ala  
 355 360 365  
 Lys Asp Lys Val Ser Leu Thr Cys Met Ile Thr Asp Phe Phe Pro Glu  
 370 375 380  
 Asp Ile Thr Val Glu Trp Gln Trp Asn Gly Gln Pro Ala Glu Asn Tyr  
 385 390 395 400  
 Lys Asn Thr Gln Pro Ile Met Asp Thr Asp Gly Ser Tyr Phe Val Tyr  
 405 410 415  
 Ser Lys Leu Asn Val Gln Lys Ser Asn Trp Glu Ala Gly Asn Thr Phe  
 420 425 430  
 Ile Cys Ser Val Leu His Glu Phe Gly Pro Glu Thr Leu Cys Gly Ala  
 435 440 445  
 Glu Leu Val Asp Ala Leu Gln Phe Val Cys Gly Asp Arg Gly Phe Tyr  
 450 455 460  
 Phe Asn Lys Pro Thr Gly Tyr Gly Ser Ser Ser Arg Arg Ala Pro Gln  
 465 470 475 480  
 Thr Gly Ile Val Asp Glu Cys Cys Phe Arg Ser Cys Asp Leu Arg Arg  
 485 490 495  
 Leu Glu Met Tyr Cys Ala Pro Leu Lys Pro Ala Lys Ser Ala  
 500 505

&lt;210&gt; 13



&lt;211&gt; 493

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Fusion protein with N-terminal portion of DAV-1 heavy chain  
and EGF mature peptide

&lt;400&gt; 13

```

Met Gly Trp Ser Trp Ile Phe Leu Phe Leu Leu Ser Gly Thr Ala Gly
 1      5      10      15
Val His Ser Glu Val Gln Leu Gln Ser Gly Pro Glu Leu Val Lys
 20      25      30
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe
 35      40      45
Thr Asp Tyr Asn Met His Trp Val Lys Gln Ser His Gly Lys Ser Leu
 50      55      60
Glu Trp Ile Gly Tyr Ile Tyr Pro Tyr Lys Gly Gly Thr Gly Tyr Asn
 65      70      75      80
Gln Lys Phe Lys Ser Lys Ala Thr Leu Thr Thr Asp Ser Ser Ser Asn
 85      90      95
Thr Ala Tyr Met Glu Leu Arg Ser Leu Thr Ser Asp Ala Ser Ala Val
 100     105     110
Tyr Tyr Cys Ala Arg Gly Ile Ala Tyr Trp Gly Gln Gly Thr Leu Val
 115     120     125
Thr Val Ser Ala Ala Lys Thr Thr Pro Pro Ser Val Tyr Pro Leu Ala
 130     135     140
Pro Gly Ser Ala Ala Gln Thr Asn Ser Met Val Thr Leu Gly Cys Leu
 145     150     155     160
Val Lys Gly Tyr Phe Pro Glu Pro Val Thr Val Thr Trp Asn Ser Gly
 165     170     175
Ser Leu Ser Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Asp
 180     185     190
Leu Tyr Thr Leu Ser Ser Ser Val Thr Val Pro Ser Ser Thr Trp Pro
 195     200     205
Ser Glu Thr Val Thr Cys Asn Val Ala His Pro Ala Ser Ser Thr Lys
 210     215     220
Val Asp Lys Lys Ile Val Pro Arg Asp Cys Gly Cys Lys Pro Cys Ile
 225     230     235     240
Cys Thr Val Pro Glu Val Ser Ser Val Phe Ile Phe Pro Pro Lys Pro
 245     250     255
Lys Asp Val Leu Thr Ile Thr Leu Thr Pro Lys Val Thr Cys Val Val
 260     265     270
Val Asp Ile Ser Lys Asp Asp Pro Glu Val Gln Phe Ser Trp Phe Val
 275     280     285
Asp Asp Val Glu Val His Thr Ala Gln Thr Gln Pro Arg Glu Glu Gln
 290     295     300
Phe Asn Ser Thr Phe Arg Ser Val Ser Glu Leu Pro Ile Met His Gln
 305     310     315     320
Asp Trp Leu Asn Gly Lys Glu Phe Lys Cys Arg Val Asn Ser Ala Ala
 325     330     335
Phe Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Arg Pro
 340     345     350
Lys Ala Pro Gln Val Tyr Thr Ile Pro Pro Pro Lys Glu Gln Met Ala
 355     360     365
Lys Asp Lys Val Ser Leu Thr Cys Met Ile Thr Asp Phe Phe Pro Glu
 370     375     380
Asp Ile Thr Val Glu Trp Gln Trp Asn Gly Gln Pro Ala Glu Asn Tyr
 385     390     395     400
Lys Asn Thr Gln Pro Ile Met Asp Thr Asp Gly Ser Tyr Phe Val Tyr
 405     410     415
Ser Lys Leu Asn Val Gln Lys Ser Asn Trp Glu Ala Gly Asn Thr Phe
 420     425     430
Ile Cys Ser Val Leu His Glu Phe Asn Ser Asp Ser Glu Cys Pro Leu
 435     440     445
Ser His Asp Gly Tyr Cys Leu His Asp Gly Val Cys Met Tyr Ile Glu
 450     455     460

```

Ala Leu Asp Lys Tyr Ala Cys Asn Cys Val Val Gly Tyr Ile Gly Glu  
 465 470 475 480  
 Arg Cys Gln Tyr Arg Asp Leu Lys Trp Trp Glu Leu Arg  
 485 490

<210> 14  
 <211> 613  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Fusion protein with N-terminal portion of DAV-1 heavy chain  
 and SCF mature peptide

<400> 14  
 Met Gly Trp Ser Trp Ile Phe Leu Phe Leu Leu Ser Gly Thr Ala Gly  
 1 5 10 15  
 Val His Ser Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys  
 20 25 30  
 Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 35 40 45  
 Thr Asp Tyr Asn Met His Trp Val Lys Gln Ser His Gly Lys Ser Leu  
 50 55 60  
 Glu Trp Ile Gly Tyr Ile Tyr Pro Tyr Lys Gly Thr Gly Tyr Asn  
 65 70 75 80  
 Gln Lys Phe Lys Ser Lys Ala Thr Leu Thr Thr Asp Ser Ser Ser Asn  
 85 90 95  
 Thr Ala Tyr Met Glu Leu Arg Ser Leu Thr Ser Asp Ala Ser Ala Val  
 100 105 110  
 Tyr Tyr Cys Ala Arg Gly Ile Ala Tyr Trp Gly Gln Gly Thr Leu Val  
 115 120 125  
 Thr Val Ser Ala Ala Lys Thr Thr Pro Pro Ser Val Tyr Pro Leu Ala  
 130 135 140  
 Pro Gly Ser Ala Ala Gln Thr Asn Ser Met Val Thr Leu Gly Cys Leu  
 145 150 155 160  
 Val Lys Gly Tyr Phe Pro Glu Pro Val Thr Val Thr Trp Asn Ser Gly  
 165 170 175  
 Ser Leu Ser Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Asp  
 180 185 190  
 Leu Tyr Thr Leu Ser Ser Ser Val Thr Val Pro Ser Ser Thr Trp Pro  
 195 200 205  
 Ser Glu Thr Val Thr Cys Asn Val Ala His Pro Ala Ser Ser Thr Lys  
 210 215 220  
 Val Asp Lys Lys Ile Val Pro Arg Asp Cys Gly Cys Lys Pro Cys Ile  
 225 230 235 240  
 Cys Thr Val Pro Glu Val Ser Ser Val Phe Ile Phe Pro Pro Lys Pro  
 245 250 255  
 Lys Asp Val Leu Thr Ile Thr Leu Thr Pro Lys Val Thr Cys Val Val  
 260 265 270  
 Val Asp Ile Ser Lys Asp Asp Pro Glu Val Gln Phe Ser Trp Phe Val  
 275 280 285  
 Asp Asp Val Glu Val His Thr Ala Gln Thr Gln Pro Arg Glu Glu Gln  
 290 295 300  
 Phe Asn Ser Thr Phe Arg Ser Val Ser Glu Leu Pro Ile Met His Gln  
 305 310 315 320  
 Asp Trp Leu Asn Gly Lys Glu Phe Lys Cys Arg Val Asn Ser Ala Ala  
 325 330 335  
 Phe Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr Lys Gly Arg Pro  
 340 345 350  
 Lys Ala Pro Gln Val Tyr Thr Ile Pro Pro Pro Lys Glu Gln Met Ala  
 355 360 365  
 Lys Asp Lys Val Ser Leu Thr Cys Met Ile Thr Asp Phe Phe Pro Glu  
 370 375 380  
 Asp Ile Thr Val Glu Trp Gln Trp Asn Gly Gln Pro Ala Glu Asn Tyr  
 385 390 395 400  
 Lys Asn Thr Gln Pro Ile Met Asp Thr Asp Gly Ser Tyr Phe Val Tyr  
 405 410 415

Ser Lys Leu Asn Val Gln Lys Ser Asn Trp Glu Ala Gly Asn Thr Phe  
 420 425 430  
 Ile Cys Ser Val Leu His Glu Phe Cys Arg Tyr Pro Ala Gln Trp Arg  
 435 440 445  
 Pro Gln Gly Ile Cys Arg Asn Arg Val Thr Asn Asn Val Lys Asp Val  
 450 455 460  
 Thr Lys Leu Val Ala Asn Leu Pro Lys Asp Tyr Met Ile Thr Leu Lys  
 465 470 475 480  
 Tyr Val Pro Gly Met Asp Val Leu Pro Ser His Cys Trp Ile Ser Glu  
 485 490 495  
 Met Val Val Gln Leu Ser Asp Ser Leu Thr Asp Leu Leu Asp Lys Phe  
 500 505 510  
 Ser Asn Ile Ser Glu Gly Leu Ser Asn Tyr Ser Ile Ile Asp Lys Leu  
 515 520 525  
 Val Asn Ile Val Asp Asp Leu Val Glu Cys Val Lys Glu Asn Ser Ser  
 530 535 540  
 Lys Asp Leu Lys Lys Ser Phe Lys Ser Pro Glu Pro Arg Leu Phe Thr  
 545 550 555 560  
 Pro Glu Glu Phe Phe Arg Ile Phe Asn Arg Ser Ile Asp Ala Phe Lys  
 565 570 575  
 Asp Phe Val Val Ala Ser Glu Thr Ser Asp Cys Val Val Ser Ser Thr  
 580 585 590  
 Leu Ser Pro Glu Lys Asp Ser Arg Val Ser Val Thr Lys Pro Phe Met  
 595 600 605  
 Leu Pro Pro Val Ala  
 610

<210> 15  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR primer for amplification of CH3 region of  
 DAV-1 heavy chain.

<400> 15  
 cctgctctgt gtttacatga ggg

23

<210> 16  
 <211> 19  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR primer for amplification of CH1 region of  
 DAV-1 heavy chain.

<400> 16  
 cccagggtca tggagtttag

19

<210> 17  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR primer for amplification of DAV-1 kappa chain  
 CL-A.

<400> 17  
 aagatggata cagttggtgc

20

<210> 18  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR primer for amplification of DAV-1 kappa chain  
 CL-B.

<400> 18  
 tgtcaagagc ttcaacagga 20

<210> 19  
 <211> 15  
 <212> PRT  
 <213> Adenovirus

<220>  
 <221> PEPTIDE  
 <222> (0)...(0)  
 <223> Peptide spanning integrin binding site on penton base.

<400> 19  
 Met Asn Asp His Ala Ile Arg Gly Asp Thr Phe Ala Thr Arg Ala  
 1 5 10 15

<210> 20  
 <211> 9  
 <212> PRT  
 <213> Adenovirus

<220>  
 <221> PEPTIDE  
 <222> (0)...(0)  
 <223> Epitope on penton base integrin binding site recognized by DAV-1.

<400> 20  
 Ile Arg Gly Asp Thr Phe Ala Thr Arg  
 1 5

<210> 21  
 <211> 31  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR sense primer for subcloning DAV-1 heavy chain for whole antibody  
 or Fab'2 constructs.

<400> 21  
 ggtaccgcca ccatgggatg gagctggatc t 31

<210> 22  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR antisense primer for subcloning DAV-1 heavy chain for  
 whole antibody construct.

<400> 22  
 gaattcatgt aacacagagc agga 24

<210> 23  
 <211> 35  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR sense primer for subcloning DAV-1 light chain for  
 whole antibody or Fab'2 constructs.

<400> 23  
aagcttgcca ccatggagac agacacaatc ctgct 35

<210> 24  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR antisense primer for subcloning DAV-1 light chain for whole antibody or Fab'2 constructs.

<400> 24  
tctagatgtc tctaacactc attcctgt 28

<210> 25  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR antisense primer for subcloning DAV-1 heavy chain for Fab'2 constructs.

<400> 25  
gaattctgat acttctggga ctgt 24

<210> 26  
<211> 26  
<212> DNA  
<213> Artificial Sequence.

<220>  
<223> PCR sense primer for subcloning TNF $\alpha$  into DAV-1/TNF $\alpha$  fusion construct.

<400> 26  
gaattcgtca gatcatcttc tcgaac 26

<210> 27  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR antisense primer for subcloning TNF $\alpha$  into DAV-1/TNF $\alpha$  fusion construct.

<400> 27  
gaattctaca gggcaatgat cccaaa 26

<210> 28  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR sense primer for subcloning IGF-1 into DAV-1/IGF-1 fusion construct.

<400> 28  
gaattcggac cggagacgct ctgcgg 26

<210> 29  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR antisense primer for subcloning IGF-1 into DAV-1/IGF-1 fusion construct.

<400> 29  
gaattctaag ctgacttggc aggctt 26

<210> 30  
<211> 96  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR sense primer for subcloning EGF into DAV-1/EGF fusion construct.

<400> 30  
gaattcaata gtgactctga atgtccctg tcccacgatg ggtactgcct ccatgatggt 60  
gtgtgcatgt atattgaagc attggacaag tatgca 96

<210> 31  
<211> 98  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR antisense primer for subcloning EGF into DAV-1/EGF fusion construct.

<400> 31  
gaattctagc gcagttccca ccacttcagg tctcgggtact gacatcgctc cccgatgtag 60  
ccaacaacac agttgcatgc atacttgtcc aatgcttc 98

<210> 32  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR sense primer for subcloning SCF into DAV-1/SCF fusion construct.

<400> 32  
gcggccgcaa gggatctgca ggaatcg 27

<210> 33  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR antisense primer for subcloning SCF into DAV-1/SCF fusion construct.

<400> 33  
tctagagtgc aacagggggt aacata 26